arXiv:math/0605205v2 [math.CV] 15 May 2006

On Uniform Subalgebras of L^∞ on the Unit Circle Generated by Almost Periodic Functions

Alexander Brudnyi*
Department of Mathematics and Statistics
University of Calgary, Calgary
Canada

Damir Kinzebulatov
Department of Mathematics and Statistics
University of Calgary, Calgary
Canada

Abstract

In the present paper we introduce analogs of almost periodic functions for the unit circle. We study certain uniform algebras generated by such functions, prove corona theorems for them and describe their maximal ideal spaces.

1. Formulation of Main Results

1.1. The classical almost periodic functions on the real line as first introduced by H. Bohr in the 1920s play an important role in various areas of Analysis. In the present paper we define analogs of almost periodic functions on the unit circle. We study certain uniform algebras generated by such functions. In particular, we describe in these terms some uniform subalgebras of the algebra H^∞ of bounded holomorphic functions on the open unit disk $\mathbb{D} \subset \mathbb{C}$, having, in a sense, the weakest possible discontinuities on the boundary $\partial\mathbb{D}$.

To formulate the main results of the paper we first recall the definition of almost periodic functions, see [B].

---

*Research supported in part by NSERC.
2000 *Mathematics Subject Classification.* Primary 30H05, Secondary 46J20.
Key words and phrases. Bounded holomorphic function, almost periodic function, uniform algebra, maximal ideal space, corona theorem.

Definition 1.1 *A continuous function* $f : \mathbb{R} \to \mathbb{C}$ *is called almost periodic if, for any* $\epsilon > 0$, *there exists* $l(\epsilon) > 0$ *such that for every* $t_0 \in \mathbb{R}$ *the interval* $[t_0, t_0 + l(\epsilon)]$ *contains at least one number* τ *for which*

$$|f(t) - f(t+\tau)| < \epsilon \quad \text{for all} \quad t \in \mathbb{R}.$$

It is well known that every almost periodic function f is uniformly continuous and is the uniform limit of a sequence of exponential polynomials $\{q_n\}_{n\in\mathbb{N}}$ where $q_n(t) := \sum_{k=1}^{n} c_{kn} e^{i\lambda_{kn} t}$, $c_{kn} \in \mathbb{C}$, $\lambda_{kn} \in \mathbb{R}$, $1 \le k \le n$, and $i := \sqrt{-1}$.

In what follows we consider $\partial\mathbb{D}$ with the *counterclockwise orientation.* For $t_0 \in \mathbb{R}$ let $\gamma_{t_0^k}(s) := \{e^{i(t_0+kt)} \ : \ 0 < t < s \le 2\pi\} \subset \partial\mathbb{D}$, $k \in \{-1, 1\}$, be two open arcs having e^{it_0} as the right or the left endpoints with respect to the chosen orientation, respectively.

Let us define almost periodic functions on open arcs of $\partial\mathbb{D}$.

Definition 1.2 *A continuous function* $f_k : \gamma_{t_0^k}(s) \to \mathbb{C}$, $k \in \{-1,1\}$, *is said to be almost periodic if the function* $\widehat{f_k} : (-\infty, 0) \to \mathbb{C}$, $\widehat{f_k}(t) := f_k(e^{i(t_0+kse^t)})$, *is the restriction of an almost periodic function on* $\mathbb{R}$.

Example 1.3 The function $e^{i\lambda \log_{t_0^k}}$, $\lambda \in \mathbb{R}$, where

$$\log_{t_0^k}(e^{i(t_0+kt)}) := \ln t, \quad 0 < t < 2\pi, \quad k \in \{-1, 1\},$$

is almost periodic in the sense of this definition on $\gamma_{t_0^1}(2\pi) = \gamma_{t_0^{-1}}(2\pi)$.

By $AP(\partial\mathbb{D}) \subset L^\infty(\partial\mathbb{D})$ we denote the uniform subalgebra of functions f such that for each t_0 and any $\epsilon > 0$ there are a number $s := s(t_0, \epsilon) \in (0, \pi)$ and almost periodic functions $f_k : \gamma_{t_0^k}(s) \to \mathbb{C}$, $k \in \{-1, 1\}$, such that

$$\operatorname*{esssup}_{z\in\gamma_{t_0^1}(s)} |f(z) - f_1(z)| < \epsilon \quad \text{and} \quad \operatorname*{ess\,sup}_{z\in\gamma_{t_0^{-1}}(s)} |f(z) - f_{-1}(z)| < \epsilon. \tag{1.1}$$

Let $S \subset \partial\mathbb{D}$ be a nonempty closed subset. By $AP(S) \subset AP(\partial\mathbb{D})$ we denote the uniform algebra of functions from $AP(\partial\mathbb{D})$ continuous on $\partial\mathbb{D} \setminus S$.

Fix a real continuous function g on $\gamma_{t_0^1}(2\pi)$ such that

$$\lim_{t\to 0+} g(e^{it}) = 1, \qquad \lim_{t\to 2\pi-} g(e^{it}) = 0$$

and $g(e^{it})$ is decreasing for $0 < t < 2\pi$. We set $g_{t_0}(e^{it}) := g(e^{i(t_0+t)})$.

Theorem 1.4 *The algebra* $AP(S)$ *is the uniform closure in* $L^\infty(\partial\mathbb{D})$ *of the algebra of complex polynomials in variables* g_{t_0} *and* $e^{i\lambda \log_{t_0^k}}$, $\lambda \in \mathbb{R}$, $e^{it_0} \in S$, $k \in \{-1, 1\}$.

Let $\phi : \partial\mathbb{D} \to \partial\mathbb{D}$ be a C^1 diffeomorphism. By $\phi^* : C(\partial\mathbb{D}) \to C(\partial\mathbb{D})$, $\phi^*(f) := f \circ \phi$, we denote the pullback by ϕ. Set $\widetilde{S} := \phi(S)$. As a consequence of Theorem 1.4 we obtain

Corollary 1.5 *ϕ^* maps $AP(\widetilde{S})$ isomorphically onto $AP(S)$.*

1.2. We say that a complex-valued function $g \in L^\infty(\partial\mathbb{D})$ has a discontinuity of the *first kind* at x_0, if the one-sided limits of g at x_0 exist but have distinct values. For a closed subset $S \subset \partial\mathbb{D}$ by $R_S \subset L^\infty(\partial\mathbb{D})$ we denote the uniform algebra of complex functions allowing discontinuities of the first kind at points of S and continuous on $\partial\mathbb{D} \setminus S$. Elements from R_S are often referred to as *regulated functions* [D]. Clearly, $R_S \hookrightarrow AP(S)$. Also, we will show (see Lemma 3.1 below) that R_S is the uniform closure of the algebra generated by all possible subalgebras R_F with finite $F \subset S$.

Let $\mathcal{M}(AP(S))$ be the *maximal ideal space* of $AP(S)$, that is, the space of all *characters* (= nonzero homomorphisms $AP(S) \to \mathbb{C}$) on $AP(S)$ equipped with the weak*-topology (also known as the *Gelfand topology*) inherited from $(AP(S))^*$. By definition, $\mathcal{M}(AP(S))$ is a compact Hausdorff space. The main result formulated in this section describes the topological structure of $\mathcal{M}(AP(S))$.

Let us consider continuous embeddings of uniform algebras

$$C(\partial\mathbb{D}) \hookrightarrow R_S \hookrightarrow AP(S).$$

The dual maps to these embeddings determine continuous surjective maps of the corresponding maximal ideal spaces:

$$\mathcal{M}(AP(S)) \xrightarrow{r_S} \mathcal{M}(R_S) \xrightarrow{c_S} \mathcal{M}(C(\partial\mathbb{D})) \cong \partial\mathbb{D}.$$

Theorem 1.6 *(1) For each $z \in S$ preimage $c_S^{-1}(z)$ consists of two points z_+ and z_- which are naturally identified with counterclockwise and clockwise orientations of $\partial\mathbb{D}$ at z.*

(2) $c_S : \mathcal{M}(R_S) \setminus c_S^{-1}(S) \to \partial\mathbb{D} \setminus S$ is a homeomorphism.



Figure 1. *For a given* $S = \{z_1, z_2, z_3, z_4\}$ *we have the homeomorphism* c_S *of*

$$\mathcal{M}(R_S) \setminus \{z_{1+}, z_{1-}, z_{2+}, z_{2-}, z_{3+}, z_{3-}, z_{4+}, z_{4-}\}$$

and $\partial\mathbb{D} \setminus \{z_1, z_2, z_3, z_4\}$ *where* $c_S(z_{i+}) = c_S(z_{i-}) = z_i$.

(3) $c_S^{-1}(S) \subset \mathcal{M}(R_S)$ *is a totally disconnected compact Hausdorff space.*

(4) For each $\xi \in c_S^{-1}(S)$ *preimage* $r_S^{-1}(\xi)$ *is homeomorphic to the Bohr compactification* $b\mathbb{R}$ *of* $\mathbb{R}$.

(5) The map $r_S : \mathcal{M}(AP(S)) \setminus (c_S \circ r_S)^{-1}(S) \to \mathcal{M}(R_S) \setminus c_S^{-1}(S) \cong \partial\mathbb{D} \setminus S$ *is a homeomorphism.*



Figure 2. *Given an* n *point set* S, *the maximal ideal space* $\mathcal{M}(AP(S))$ *is the union of* $\partial\mathbb{D} \setminus S$ *and* $2n$ *Bohr compactifications of* $\mathbb{R}$ *that can be viewed as infinite dimensional "tori", where the spirals joining the arcs and the tori are meant to indicate (in a figurative manner) that there is an influence of the topology of the Bohr compactifications on the topology of the arcs.*

Let us recall that $b\mathbb{R}$ is a compact abelian topological group homeomorphic to the maximal ideal space of the algebra of continuous almost periodic functions on $\mathbb{R}$. Also, it follows straightforwardly from (2)-(5) that

(6) The covering dimension of $\mathcal{M}(AP(S))$ is ∞.

(7) For a continuous map $\phi : T \to (c_S \circ r_S)^{-1}(S)$ of a connected topological space T, there is a point $\xi \in c_S^{-1}(S)$ such that $\phi(T) \subset r_S^{-1}(\xi)$.

1.3. By $A_0 \subset H^\infty$ we denote the *disk-algebra*, i.e., the algebra of functions continuous on the closure $\overline{\mathbb{D}}$ and holomorphic in $\mathbb{D}$. Also, by $f|_{\partial\mathbb{D}}$ we denote the boundary values of $f \in C(\mathbb{D})$ (in case they exist). In the present part we describe uniform subalgebras of H^∞ generated by almost periodic functions. These subalgebras contain A_0 and have, in a sense, the weakest possible discontinuities on $\partial\mathbb{D}$.

Suppose that S contains at least 2 points. By $A_S \subset H^\infty$ we denote the uniform closure of the algebra generated by A_0 and by holomorphic functions of the form e^f where $\mathrm{Re}f|_{\partial\mathbb{D}}$ is a finite linear combination with real coefficients of characteristic functions of closed arcs whose endpoints belong to S. If S consists of a single point we determine $A_S \subset H^\infty$ to be the uniform closure of the algebra generated by A_0 and functions $ge^{\lambda f}$, $\lambda \in \mathbb{R}$, where $\mathrm{Re}f|_{\partial\mathbb{D}}$ is the characteristic function of a closed arc with an endpoint at S and $g \in A_0$ is a function such that ge^f has discontinuity on S only. In the following result we naturally identify A_S and H^∞ with the algebras of their boundary values.

Theorem 1.7

$$A_S = AP(S) \cap H^\infty.$$

Remark 1.8 Suppose that $F \subset \partial\mathbb{D}$ contains at least 2 points. Let $e^{\lambda f} \in A_S$, $\lambda \in \mathbb{R}$, where $\mathrm{Re}\, f$ is the characteristic function of an arc $[x, y]$ with $x, y \in S$. Let $\phi_{x,y} : \mathbb{D} \to \mathbb{H}_+$ be the bilinear map onto the upper half-plane that maps x to 0, the midpoint of the arc $[x, y]$ to 1 and y to ∞. Then there is a constant C such that

$$e^{\lambda f(z)} = e^{-\frac{i\lambda}{\pi}\mathrm{Log}\,\phi_{x,y}(z)+\lambda C}, \quad z \in \mathbb{D},$$

where Log denotes the principal branch of the logarithmic function. Thus from Theorem 1.7 it follows that the algebra $AP(S) \cap H^\infty$ is the uniform closure of the algebra generated by A_0 and the functions $e^{i\lambda(\mathrm{Log}\circ\phi_{x,y})}$, $\lambda \in \mathbb{R}$, $x, y \in S$.

The following example shows that if S is an infinite set, A_S does not coincide with the algebra generated by functions e^f where $\mathrm{Re}f \in R_S$ (the corresponding arguments are presented in section 4.3).

Example 1.9 Assume that a closed subset $S \subset \partial\mathbb{D}$ contains -1, 1 and a sequence $\{e^{it_k}\}_{k\in\mathbb{N}}$, $t_k \in (0, \pi/2)$, converging to 1. Let $\{\alpha_k\}_{k\in\mathbb{N}}$ be a sequence of positive numbers satisfying the condition $\sum_{k=1}^{n} \alpha_k = 1$. By χ_k we denote the characteristic function of the arc $\gamma_k := \{e^{it} \ : \ t_k \le t \le \pi\}$. Consider the function

$$u(z) := \sum_{k=1}^{n} \alpha_k \chi_k(z), \quad z \in \partial\mathbb{D}.$$

Clearly, $u \in R_S$. Let h be a holomorphic function on $\mathbb{D}$ such that $\mathrm{Re}h|_{\partial\mathbb{D}} = u$. Then $e^h \in H^\infty \setminus A_S$. However, for any $f \in A_0$ such that $f(1) = 0$ we have $fe^h \in A_S$.

Remark 1.10 It seems to be natural that a function $f \in H^\infty$ has the weakest possible discontinuities on $\partial\mathbb{D}$ if $f|_{\partial\mathbb{D}} \in R_S$. However, from the classical Lindelöf theorem [L] it follows that any such f in fact belongs to A_0. Moreover, the same conclusion is obtained even from the fact that $\mathrm{Re}f|_{\partial\mathbb{D}} \in R_S$ for $f \in H^\infty$. In particular, if f is holomorphic on $\mathbb{D}$ and $\mathrm{Re}f|_{\partial\mathbb{D}}$ is correctly defined and belongs to $R_S \setminus C(\partial\mathbb{D})$, then $f \notin H^\infty$.[1] Nevertheless, $e^f \in H^\infty$, which partly explains the choice of the object of our research.

Let $\mathcal{M}(A_S)$ be the maximal ideal space of A_S. Since evaluation functionals $z(f) := f(z)$, $z \in \mathbb{D}$, $f \in A_S$, belong to $\mathcal{M}(A_S)$ and A_S separates points on $\mathbb{D}$, there is a continuous embedding $i_S : \mathbb{D} \to \mathcal{M}(A_S)$. In the sequel we identify $\mathbb{D}$ with $i_S(\mathbb{D})$. Then the following corona theorem is true.

Theorem 1.11 *$\mathbb{D}$ is dense in $\mathcal{M}(A_S)$.*

Remark 1.12 Let us recall that the corona theorem is equivalent to the following statement, see, e.g., [G, Chapter V]:

For any collection of functions $f_1, \dots, f_n \in A_S$ satisfying the corona condition

$$\max_{1\le j\le n} |f_j(z)| \ge \delta > 0, \quad z \in \mathbb{D}, \tag{1.2}$$

there are functions $g_1, \dots, g_n \in A_S$ such that

$$f_1 g_1 + \dots + f_n g_n = 1. \tag{1.3}$$

Finally we formulate some results on the structure of $\mathcal{M}(A_S)$. Since $A_0 \hookrightarrow A_S$, there is a continuous surjection of the maximal ideal spaces

$$a_S : \mathcal{M}(A_S) \to \mathcal{M}(A_0) \cong \overline{\mathbb{D}}.$$

Recall that the *Šilov boundary* of A_S is the smallest compact subset $K \subset \mathcal{M}(A_S)$ such that for each $f \in A_S$

$$\sup_{z\in\mathcal{M}(A_S)} |f(z)| = \sup_{\xi\in K} |f(\xi)|.$$

Here we assume that every $f \in A_S$ is also defined on $\mathcal{M}(A_S)$ where its extension to $\mathcal{M}(A_S) \setminus \mathbb{D}$ is given by the *Gelfand transform*: $f(\xi) := \xi(f)$, $\xi \in \mathcal{M}(A_S)$.

Theorem 1.13 *(1) $a_S : \mathcal{M}(A_S) \setminus a_S^{-1}(S) \to \overline{\mathbb{D}} \setminus S$ is a homeomorphism.*

(2) The Šilov boundary K_S of A_S is naturally homeomorphic to $\mathcal{M}(AP(S))$. Under the identification of K_S and $\mathcal{M}(AP(S))$ one has $a_S|_{K_S} = r_S \circ c_S$.

---

[1] In this case $f|_{\partial\mathbb{D}} \in BMO(\partial\mathbb{D})$ with $||f||_{BMO(\partial\mathbb{D})} \le c||\mathrm{Re}f|_{\partial\mathbb{D}}||_{L^\infty(\partial\mathbb{D})}$ for some absolute constant $c > 0$.

(3) For each $z \in S$, the preimage $a_S^{-1}(z)$ is homeomorphic to the maximal ideal space of the algebra $AP_{\mathcal{O}}(\Sigma)$ of uniformly continuous almost periodic functions on the strip $\Sigma := \{z \in \mathbb{C} \ : \ \operatorname{Im} z \in [0, \pi]\}$ holomorphic at interior points of Σ.

In the next section we describe the topological structure of the maximal ideal space $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ of $AP_{\mathcal{O}}(\Sigma)$. We show that this space is equipped with a natural "complex structure". Similarly each fibre $a_S^{-1}(z)$, $z \in S$, has a natural complex structure so that the homeomorphisms in Theorem 1.13 (3) are analytic with respect to these structures.

In a forthcoming paper we present similar results for algebras of bounded holomorphic functions on open polydisks generated by almost periodic functions.

2. Maximal Ideal Space of the Algebra $AP_{\mathcal{O}}(\Sigma)$

2.1. The construction presented below is rather general and can be defined for Galois coverings of complex manifolds with boundaries (cf. [Br]). However, we restrict ourselves to the case of coverings of annuli related to the subject of our paper.

Consider the annulus $R := \{z \in \mathbb{C} \ : \ e^{-2\pi^2} \leq |z| \leq 1\}$. Its universal covering can be identified with Σ so that $e : \Sigma \to R$, $e(z) := e^{2\pi i z}$, $z \in \Sigma$, is the covering map. We can also regard Σ as a principal bundle on R with fibre $\mathbb{Z}$ (see, e.g., [H] for the corresponding topological definitions). To specify, consider a cover of R by relatively open simply connected sets U_1 and U_2. Then $e^{-1}(U_k)$ can be identified with $U_k \times \mathbb{Z}$, $k = 1, 2$. Also, there is a continuous map $c_{12} : U_1 \cap U_2 \to \mathbb{Z}$ such that R is isomorphic (in the category of complex manifolds with boundaries) to the quotient space of $(U_1 \times \mathbb{Z}) \sqcup (U_2 \times \mathbb{Z})$ under the equivalence relation:

$$U_1 \times \mathbb{Z} \ni (z, n) \sim (z, n + c_{12}(z)) \in U_2 \times \mathbb{Z} \text{ for all } z \in U_1 \cap U_2 \text{ and } n \in \mathbb{Z}.$$

Let $b\mathbb{Z}$ be the Bohr compactification of $\mathbb{Z}$. Then the action of $\mathbb{Z}$ on itself by translations can be extended naturally to the action on $b\mathbb{Z}$: $\xi \mapsto \xi + n$, $\xi \in b\mathbb{Z}$, $n \in \mathbb{Z}$. By $E(R, b\mathbb{Z})$ we denote the principal bundle on R with fibre $b\mathbb{Z}$ defined as the quotient of $(U_1 \times b\mathbb{Z}) \sqcup (U_2 \times b\mathbb{Z})$ under the equivalence relation:

$$U_1 \times b\mathbb{Z} \ni (z, \xi) \sim (z, \xi + c_{12}(z)) \in U_2 \times b\mathbb{Z} \text{ for all } z \in U_1 \cap U_2 \text{ and } \xi \in b\mathbb{Z}.$$

Clearly $E(R, b\mathbb{Z})$ is a compact Hausdorff space in the quotient topology induced by that of $(U_1 \times b\mathbb{Z}) \sqcup (U_2 \times b\mathbb{Z})$. Also, the projection $p : E(R, b\mathbb{Z}) \to R$ is defined by the natural projections $U_k \times \mathbb{Z} \to U_k$, $k = 1, 2$, onto the first coordinate.

Next, the natural local injections $U_k \times \mathbb{Z} \hookrightarrow U_k \times b\mathbb{Z}$, $k = 1, 2$, determine an injection $i_0 : \Sigma \hookrightarrow E(R, b\mathbb{Z})$ such that $p \circ i_0 = e$. Moreover, $i_0(\Sigma)$ is dense in $E(R, b\mathbb{Z})$, because $\mathbb{Z}$ is dense in $b\mathbb{Z}$ (in the topology of $b\mathbb{Z}$). Similarly, one determines an injection $i_\xi : \Sigma \hookrightarrow E(R, b\mathbb{Z})$, $\xi \in b\mathbb{Z}$, by the formula $i_\xi((z, n)) := (z, \xi + n)$, $z \in U_k$, $n \in \mathbb{Z}$, $k = 1, 2$. Since, by definition, $\xi + \mathbb{Z}$ is dense in $b\mathbb{Z}$, the image $i_\xi(\Sigma)$ is dense in $E(R, b\mathbb{Z})$ for any ξ. Moreover, $E(R, b\mathbb{Z}) = \sqcup_\xi i_\xi(\Sigma)$ where the union is taken over all ξ whose images in the quotient group $b\mathbb{Z}/\mathbb{Z}$ are mutually distinct. Observe also that every i_ξ is a continuous map and locally is an embedding.

Definition 2.1 *A continuous function f on $E(R, b\mathbb{Z})$ is said to be holomorphic if every function $f \circ i_\xi$ is holomorphic in interior points of Σ.*

By $\mathcal{O}(E(R, b\mathbb{Z})) \subset C(E(R, b\mathbb{Z}))$ we denote the Banach algebra of holomorphic functions on $E(R, b\mathbb{Z})$.

Remark 2.2 Using a normal family argument and the fact that $i_\xi(\Sigma)$ is dense in $E(R, b\mathbb{Z})$ one can easily show that $f \in C(E(R, b\mathbb{Z}))$ is holomorphic if and only if there is $\xi \in b\mathbb{Z}$ such that $f \circ i_\xi$ is holomorphic in interior points of Σ.

2.2. Let us recall that $f \in AP_{\mathcal{O}}(\Sigma)$ if f is uniformly continuous with respect to the Euclidean metric on Σ, holomorphic in interior points of Σ and its restriction to each straight line parallel to the x-axis is almost periodic. In the following lemma we identify Σ with $i_0(\Sigma) \subset E(R, b\mathbb{Z})$.

Lemma 2.3 *Every $f \in AP_{\mathcal{O}}(\Sigma)$ admits a continuous extension to a holomorphic function $\widehat{f}$ on $E(R, b\mathbb{Z})$. Moreover, the correspondence $\widehat{\ }: AP_{\mathcal{O}}(\Sigma) \to \mathcal{O}(E(R, b\mathbb{Z}))$, $f \mapsto \widehat{f}$, determines an isomorphism of Banach algebras.*

Proof. Using the Pontryagin duality [P] one can easily show that the closure of $\mathbb{Z}$ in $b\mathbb{R}$, the Bohr compactification of $\mathbb{R}$, is isomorphic to $b\mathbb{Z}$. In particular, restrictions to $\mathbb{Z}$ of almost periodic functions on $\mathbb{R}$ are almost periodic functions on $\mathbb{Z}$ and the algebra generated by extensions of such functions to $b\mathbb{Z}$ separates points on $b\mathbb{Z}$. Let us identify naturally every $p^{-1}(U_k)$ with $U_k \times b\mathbb{Z}$, and every $e^{-1}(U_k)$ with $U_k \times \mathbb{Z}$ ($\subset U_k \times b\mathbb{Z}$), $k = 1, 2$, and regard $f|_{e^{-1}(U_k)}$, $f \in AP_{\mathcal{O}}(\Sigma)$, as a bounded function $f_k \in C(U_k \times \mathbb{Z})$. Then, f_k is

(a) holomorphic in interior points of $U_k \times \mathbb{Z}$,

(b) uniformly continuous on $U_k \times \mathbb{Z}$ with respect to the semi-metric $r(v_1, v_2) := |z_1 - z_2|$ on $U_k \times \mathbb{Z}$ where $v_1 = (z_1, n_1)$, $v_2 = (z_2, n_2) \in U_k \times \mathbb{Z}$ and $|\cdot|$ is the Euclidean norm on $\mathbb{C}$,

(c) $f_k|_{\{z\}\times\mathbb{Z}}$ is almost periodic on $\mathbb{Z}$ for every $z \in U_k$.

To prove the lemma it suffices to show that

(1) *There is a continuous function $\widehat{f_k}$ on $U_k \times b\mathbb{Z}$ such that $\widehat{f_k}|_{U_k\times\mathbb{Z}} = f_k$, for every $\xi \in b\mathbb{Z}$, the function $\widehat{f_k}|_{U_k\times\{\xi\}}$ is holomorphic in interior points of U_k and* $\sup_{U_k\times b\mathbb{Z}} |\widehat{f_k}| = \sup_{U_k\times\mathbb{Z}} |f_k|$.

(2) *If $f \in \mathcal{O}(E(R, b\mathbb{Z}))$, then $f|_\Sigma \in AP_{\mathcal{O}}(\Sigma)$.*

(1) Since for every $z \in U_k$ the function $f_{kz}(n) := f_k(z, n)$ is almost periodic on $\mathbb{Z}$, there is a continuous function $\widehat{f_{kz}}$ on $b\mathbb{Z}$ which extends f_{kz}. We set $\widehat{f_k}(z, \xi) := \widehat{f_{kz}}(\xi)$, $\xi \in b\mathbb{Z}$ and prove that $\widehat{f_k}$ is continuous. In fact, take a point $w = (z, \xi) \in U_k \times b\mathbb{Z}$ and a number $\epsilon > 0$. By the uniform continuity of f_k, there is $\delta > 0$ such that for any pair of points $v_1 = (z_1, n)$ and $v_2 = (z_2, n)$ from $U_k \times \mathbb{Z}$ with $|z_1 - z_2| < \delta$ one has $|f(z_1, n) - f(z_2, n)| < \epsilon/3$. Define a neighbourhood U_z of $z \in U_k$ by $U_z :=$

$\{z' \in U_k \ : |z - z'| < \delta\}$. Further, by the definition of $\widehat{f}_{kz}$, there is a neighbourhood $U_\xi \subset b\mathbb{Z}$ of ξ such that for any $\eta \in U_\xi$ we have $|\widehat{f}_{kz}(\eta) - \widehat{f}_{kz}(\xi)| < \epsilon/3$. Consider $U_w := U_z \times U_\xi$. Then U_w is an open neighbourhood of $w \in U_k \times b\mathbb{Z}$. Note that $f_{kz} - f_{kz'}$ is an almost periodic function on $\mathbb{Z}$ and for any $z' \in U_z$ its supremum norm is $< \epsilon/2$. This implies that $|\widehat{f}_{kz}(\eta) - \widehat{f}_{kz'}(\eta)| < \epsilon/2$ for each $\eta \in b\mathbb{Z}$. In particular, for any $(x, \eta) \in U_w$ we have

$$|\widehat{f}_k(x,\eta) - \widehat{f}_k(z,\xi)| \le |\widehat{f}_{kx}(\eta) - \widehat{f}_{kz}(\eta)| + |\widehat{f}_{kz}(\eta) - \widehat{f}_{kz}(\xi)| < \epsilon \ .$$

This shows that $\widehat{f}_k$ is continuous at every $w \in U_k \times b\mathbb{Z}$.

Now, we show that $\widehat{f}|_{U_k \times \{\xi\}}$ is holomorphic in interior points of U_k for every $\xi \in b\mathbb{Z}$.

Since $\widehat{f}_k$ is uniformly continuous on the compact set $U_k \times b\mathbb{Z}$, for any $\epsilon > 0$ there is $n_\epsilon \in \mathbb{Z}$ such that $\sup_{z \in U_k} |\widehat{f}_k(z,\xi) - f_k(z, n_\epsilon)| < \epsilon$. In particular, $\widehat{f}_k(\cdot, \xi)$ is the limit in $C(U_k)$ of the sequence $\{f_k(\cdot, n_{1/l})\}_{l \ge 1}$ of bounded continuous functions holomorphic on the interior of U_k. Thus $\widehat{f}_k|_{U_k \times \{\xi\}}$ is also holomorphic on the interior of U_k.

Note that the equality $\sup_{U_k \times b\mathbb{Z}} |\widehat{f}_k| = \sup_{U_k \times \mathbb{Z}} |f_k|$ follows directly from the definition of $\widehat{f}_k$. This completes the proof of (1).

(2) Suppose that $f \in \mathcal{O}(E(R, b\mathbb{Z}))$. We must show that $f|_\Sigma \in AP_{\mathcal{O}}(\Sigma)$. To this end it suffices to show that for every line $L := \{z \in \mathbb{C} \ : \ \operatorname{Im} z = t \in [0, \pi]\}$, the function $f|_L$ is almost periodic. (The uniform continuity of $f|_\Sigma$ with respect to the Euclidean metric on $\mathbb{C}$ follows easily from the uniform continuity of f on $E(R, b\mathbb{Z})$.) By definition the image $S := e(L) \subset R$ is a circle and $e|_L : L \to S$ is the universal covering. Consider the compact set $p^{-1}(S) \subset E(R, b\mathbb{Z})$. Since the function $f|_{p^{-1}(S)}$ is continuous and $b\mathbb{Z} \subset b\mathbb{R}$, given $\epsilon > 0$ there are a finite open cover $(V_n)_{1 \le n \le m}$ of S by sets homeomorphic to open intervals in $\mathbb{R}$ and continuous almost periodic functions f_n on L, $1 \le n \le m$, such that

$$\sup_{z \in e^{-1}(V_n), 1 \le n \le m} |f_n(z) - f(z)| < \epsilon. \tag{2.1}$$

Let $\{\rho_n\}_{1 \le n \le m}$ be a continuous partition of unity subordinate to $(V_n)_{1 \le n \le m}$. We pull back it to L by e and by $\widetilde{\rho}_n$ $(:= e^* \rho_n)$, $1 \le n \le m$, denote the obtained functions. Since each $\widetilde{\rho}_n$ is periodic on L, it is almost periodic, as well. Let us define the function f_ϵ on L by the formula

$$f_\epsilon(z) := \sum_{n=1}^{m} \widetilde{\rho}_n(z) f_n(z).$$

Then clearly f_ϵ is a continuous almost periodic function on L and by (2.1)

$$\sup_{z \in L} |f_\epsilon(z) - f(z)| < \epsilon.$$

This shows that f admits uniform approximation on L by continuous almost periodic functions and therefore $f|_L$ is almost periodic, as well.

The lemma has been proved. $\square$

2.3. In this part we prove the corona theorem for the algebra $AP_{\mathcal{O}}(\Sigma)$. Recall that $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ stands for its maximal ideal space. It is well known that every $f \in AP_{\mathcal{O}}(\Sigma)$ can be approximated uniformly on Σ by polynomials in $e^{i\lambda z}$, $\lambda \in \mathbb{R}$, see, e.g., [JT]. Then using the inverse limit construction for maximal ideal spaces of uniform algebras, see [R], one obtains that the base of topology of $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ is generated by functions $e^{i\lambda z}$. Namely, the base of topology on $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ consists of open sets of the form

$$U(\lambda_1, \dots, \lambda_l, \xi, \epsilon) := \{\eta \in \mathcal{M}(AP_{\mathcal{O}}(\Sigma)) \ : \ \max_{1\le k\le l} |e_{\lambda_k}(\eta) - e_{\lambda_k}(\xi)| < \epsilon\}$$

where e_λ is the extension of $e^{i\lambda z}$ to $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ by means of the Gelfand transform.

Theorem 2.4 *Σ is dense in $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ in the Gelfand topology.*

Proof. Assume that the corona theorem is not true for $AP_{\mathcal{O}}(\Sigma)$, that is, Σ is not dense in $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$. Then there exist $\xi \in \mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ and its neighbourhood $U(\lambda_1, \dots, \lambda_l, \xi, \epsilon)$ such that

$$U(\lambda_1, \dots, \lambda_l, \xi, \epsilon) \cap cl(\Sigma) = \emptyset;$$

here $cl(\Sigma)$ is the closure of Σ in $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$. Denoting $c_k := e_{\lambda_k}(\xi)$, $1 \le k \le l$, we have

$$\max_{1\le k\le l} |e^{i\lambda_k z} - c_k| \ge \epsilon > 0 \quad \text{for all} \quad z \in \Sigma. \tag{2.2}$$

Clearly, every function $e^{i\lambda_k z} - c_k$, $1 \le k \le l$, has at least one zero in Σ. (For otherwise, if, say, $e^{i\lambda_k z} - c_k$ has no zeros on Σ, then the function $g_k(z) := \frac{1}{e^{i\lambda_k z} - c_k}$, $z \in \Sigma$, obviously belongs to $AP_{\mathcal{O}}(\Sigma)$ and $g_k(z)(e^{i\lambda_k z} - c_k) = 1$ for all $z \in \Sigma$, a contradiction to our assumption.) In particular, since solutions of the equation

$$e^{i\lambda_k z} = c_k, \quad \lambda_k \ne 0,$$

are given by

$$z = -\frac{i \ln |c_k|}{\lambda_k} + \frac{\operatorname{Arg} c_k + 2\pi s}{\lambda_k}, \quad s \in \mathbb{Z},$$

they all belong to Σ. Further, without loss of generality we may assume that all $\lambda_k > 0$. Indeed, if some $\lambda_k < 0$, we can replace the function $e^{i\lambda_k z} - c_k$ by $e^{-i\lambda_k z} - \frac{1}{c_k}$ (observe that $c_k \ne 0$ by the above argument) so that the new family of functions also satisfies (2.2) (possibly with a different ϵ) and extensions of these functions to $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ vanish at ξ. Since all these functions have zeros in Σ and satisfy (2.2) there, we have

$$\max_{1\le k\le l} |e^{i\lambda_k z} - c_k| \ge \widetilde{\epsilon} > 0, \quad \text{for all} \quad z \in \mathbb{H}_+$$

where $\mathbb{H}_+ \subset \mathbb{C}$ is the open upper half-plane, and all $e^{i\lambda_k z} - c_k$, $\lambda_k \in \mathbb{R}_+$, are almost periodic on $\overline{\mathbb{H}}_+$. From the last inequality by the Bötcher corona theorem [Bö] we obtain that there exist holomorphic almost periodic functions $g_1, \dots, g_l$ on $\overline{\mathbb{H}}_+$ such that

$$\sum_{k=1}^{l} g_k(z)(e^{i\lambda_k z} - c_k) = 1 \quad \text{for all} \quad z \in \overline{\mathbb{H}}_+.$$

Thus taking the restrictions of these functions to Σ we obtain a contradiction to our assumption.

This completes the proof of the corona theorem for $AP_{\mathcal{O}}(\Sigma)$. $\square$

Corollary 2.5 *$E(R, b\mathbb{Z})$ is homeomorphic to $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$.*

Proof. By Lemma 2.3 there exists a continuous embedding i of $E(R, b\mathbb{Z})$ into $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$. Since Σ is dense in $E(R, b\mathbb{Z})$ and $i(\Sigma)$ is dense in $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ by the corona theorem, i is a homeomorphism. $\square$

Remark 2.6 (1) According to our construction, the closure of $\mathbb{R}$ in $E(R, b\mathbb{Z})$ coincides with $b\mathbb{R}$. In fact, this closure is $E(\partial\mathbb{D}, b\mathbb{Z})$, the principal bundle on $\partial\mathbb{D}$ with fibre $b\mathbb{Z}$ obtained as the restriction of $E(R, b\mathbb{Z})$ to $\partial\mathbb{D}$. Since R is homotopically equivalent to $\partial\mathbb{D}$, by the covering homotopy theorem $E(R, b\mathbb{Z})$ is homotopically equivalent to $b\mathbb{R} = E(\partial\mathbb{D}, b\mathbb{Z})$.
(2) Observe also that the covering dimension of $b\mathbb{R}$ is ∞, because $b\mathbb{R}$ is the inverse limit of real tori whose dimensions go to ∞. Therefore the covering dimension of $\mathcal{M}(AP_{\mathcal{O}}(\Sigma)) = E(R, b\mathbb{Z})$ is also ∞.
(3) Finally, it is easy to show that the Šilov boundary of $AP_{\mathcal{O}}(\Sigma)$ is $E(R, b\mathbb{Z})|_{\partial R}$, the restriction of $E(R, b\mathbb{Z})$ to the boundary ∂R of R, and is homeomorphic to $b\mathbb{R} \sqcup b\mathbb{R}$.

3. Proofs of Theorems 1.4, 1.6 and Corollary 1.5

3.1. Proof of Theorem 1.4. Let $f \in AP(S)$. Since $\partial\mathbb{D}$ is a compact set, given $\epsilon > 0$ there are finitely many points $z_l := e^{it_l} \in \partial\mathbb{D}$, numbers $s_l \in (0, \pi)$ and almost periodic functions $f_k^l : \gamma_{t_l^k}(s_l) \to \mathbb{C}$, $k \in \{-1, 1\}$, $1 \le l \le n$, such that

$$\begin{gathered}\bigcup_{l=1}^{n} (\gamma_{t_l^{-1}}(s_l) \cup \gamma_{t_l^1}(s_l)) = \partial\mathbb{D} \setminus \{z_1, \dots, z_n\} \quad \text{and for all} \quad 1 \le l \le n \\ \operatorname*{ess\,sup}_{z \in \gamma_{t_l^1}(s_l)} |f(z) - f_1^l(z)| < \frac{\epsilon}{2}, \qquad \operatorname*{ess\,sup}_{z \in \gamma_{t_l^{-1}}(s_l)} |f(z) - f_{-1}^l(z)| < \frac{\epsilon}{2}.\end{gathered} \tag{3.1}$$

We set $U_l := \gamma_{t_l^1}(s_l) \cup \gamma_{t_l^{-1}}(s_l) \cup \{z_l\}$. Then $U = (U_l)_{l=1}^n$ is a finite open cover of $\partial\mathbb{D}$. Let $\{\rho_l\}_{l=1}^n$ be a continuous partition of unity subordinate to U such that $\operatorname{supp} \rho_l \subset\subset U_l$ and $\rho_l(z_l) = 1$, $1 \le l \le n$. Consider the functions f_l on $\partial\mathbb{D} \setminus \{z_l\}$ defined by the formulas

$$f_l(z) := \begin{cases} \rho_l(z) f_{-1}^l(z), & \text{if} \quad z \in \gamma_{t_l^{-1}}(s_l), \\ \rho_l(z) f_1^l(z), & \text{if} \quad z \in \gamma_{t_l^1}(s_l). \end{cases}$$

Since f_l is continuous outside z_l and coincides with f_{-1}^l and f_1^l in a neighbourhood of z_l, it belongs to $AP(S)$. Moreover,

$$||f - \sum_{l=1}^{n} f_l||_{L^\infty(\partial\mathbb{D})} < \frac{\epsilon}{2}. \tag{3.2}$$

Thus in order to prove the theorem it suffices to approximate every f_l by polynomials in g_{t_l} and $e^{i\lambda \log_{t_l^k}}$, $k \in \{-1, 1\}$, $\lambda \in \mathbb{R}$.

Suppose first that $z_l \notin S$. Since f is continuous outside the compact set S, we can choose the above cover U and the family of functions $\{f_k^m\}_{1\leq m\leq n}$, $k \in \{-1, 1\}$, such that in the U_l the functions f_k^l, $k \in \{-1, 1\}$, have the same limit at z_l. This implies the continuity of f_l on $\partial\mathbb{D}$. Next, consider the uniform algebra $\mathbb{C}(g_{t_l})$ over $\mathbb{C}$ generated by the function g_{t_l}. Since by our definition g_{t_l} separates points on $\partial\mathbb{D} \setminus \{z_l\}$, the maximal ideal space of $\mathbb{C}(g_{t_l})$ is homeomorphic to the closed interval $(\partial\mathbb{D} \setminus \{z_l\}) \cup \{z_{l^k}\}$, $k \in \{-1, 1\}$, with endpoints z_{l^1} and $z_{l^{-1}}$ identified with the counterclockwise and clockwise orientations at z_l. Clearly every continuous function on $\partial\mathbb{D}$ is extended to the maximal ideal space of $\mathbb{C}(g_{t_l})$ as a continuous function having the same values at z_{l^1} and $z_{l^{-1}}$. Thus by the Stone-Weierstrass theorem f_l can be uniformly approximated on $\partial\mathbb{D} \setminus \{z_l\}$ by complex polynomials in g_{t_l}.

Now, suppose that $z_l \in S$. Choose some $s \in (0, s_l)$. By $AP_{\{z_l\}}(s)$ we denote the uniform algebra of complex continuous functions on $\partial\mathbb{D} \setminus \{z_l\}$ almost periodic on the open arcs $\gamma_{t_l^k}(s)$, $k \in \{-1, 1\}$. (Since $s \in (0, \pi)$, the closures of these arcs are disjoint.) By $\mathcal{M}_{\{z_l\}}(s)$ we denote the maximal ideal space of $AP_{\{z_l\}}(s)$. Then $\partial\mathbb{D} \setminus \{z_l\}$ is dense in $\mathcal{M}_{\{z_l\}}(s)$ (in the Gelfand topology). Note that the space $\mathcal{M}_{\{z_l\}}(s)$ is constructed as follows.

Consider the Bohr compactification $b\mathbb{R}$ of $\mathbb{R}$. We identify the negative ray $\mathbb{R}_-$ in $\mathbb{R} \subset b\mathbb{R}$ with $\gamma_{t_{l^1}}(s) \subset \partial\mathbb{D}$ by means of the map $t \mapsto e^{i(t_l + se^t)}$, $t \in \mathbb{R}_-$. Similarly, consider another copy of $b\mathbb{R}$ and identify $\mathbb{R}_-(\subset \mathbb{R})$ in this copy with $\gamma_{t_{l^{-1}}}(s) \subset \partial\mathbb{D}$ by means of the map $t \mapsto e^{i(t_l - se^t)}$, $t \in \mathbb{R}_-$. On the identified sets we introduce the topology induced from $b\mathbb{R}$ and on $\partial\mathbb{D}\setminus(\gamma_{t_{l^{-1}}}(s)\cup\gamma_{t_{l^1}}(s))$ the topology is induced from $\partial\mathbb{D}$. Then the quotient space of $b\mathbb{R} \sqcup b\mathbb{R} \sqcup \partial\mathbb{D}$ under these identifications coincides with $\mathcal{M}_{\{z_l\}}(s)$.

Next, recall that by definition the algebra $AP(\{z_l\})$ is the uniform closure in $C(\partial\mathbb{D} \setminus \{z_l\})$ of the algebra generated by the algebras $AP_{\{z_l\}}(s)$, $s \in (0, s_l)$. By $\mathcal{M}(AP(\{z_l\}))$ we denote the maximal ideal space of $AP(\{z_l\})$. Since for any $s'' < s'$ we have inclusions $i_{s''s'} : AP_{\{z_l\}}(s') \hookrightarrow AP_{\{z_l\}}(s'')$, the space $\mathcal{M}(AP(\{z_l\}))$ is the inverse limit of the spaces $\mathcal{M}_{\{z_l\}}(s)$ (here the maps $p_{s''s'} : \mathcal{M}_{\{z_l\}}(s'') \to \mathcal{M}_{\{z_l\}}(s')$ in the definition of this limit are defined as the dual maps to $i_{s''s'}$). Also, $\partial\mathbb{D} \setminus \{z_l\}$ is dense in $\mathcal{M}(AP(\{z_l\}))$ in the Gelfand topology. Since by the definition the functions f_l, $e^{i\lambda \log_{t_{l^k}}}$ and g_{t_l} admit the continuous extensions (denoted by the same symbols) to $\mathcal{M}(AP(\{z_l\}))$, it suffices to show that the extended functions $e^{i\lambda \log_{t_{l^k}}}$, g_{t_l} separate points on $\mathcal{M}(AP(\{z_l\}))$. Then we will apply the Stone-Weierstrass theorem to get a complex polynomial p_l in variables $e^{i\lambda \log_{t_l^k}}$ and g_{t_l} which uniformly approximates f_l on $\mathcal{M}(AP(\{z_l\}))$ with an error $< \epsilon/2n$. Therefore, $\sum_{l=1}^n p_l$ will approximate f in $L^\infty(\partial\mathbb{D})$ with an error $< \epsilon$.

So let us show that the functions $e^{i\lambda\log_{t_l^k}}$, g_{t_l} separate points on $\mathcal{M}(AP(\{z_l\}))$. By $p_s : \mathcal{M}(AP(\{z_l\})) \to \mathcal{M}_{\{z_l\}}(s)$ we denote the continuous surjection determined by the inverse limit construction. First, we consider distinct points $x, y \in \mathcal{M}(AP(\{z_l\}))$ for which there is $s \in (0, s_l)$ such that $p_s(x)$ and $p_s(y)$ are distinct and belong to one of the Bohr compactifications of $\mathbb{R}$ in $\mathcal{M}_{\{z_l\}}(s)$, say, e.g., to the compactification obtained by gluing $\mathbb{R}_-$ with $\gamma_{t_{l1}}(s)$. Since in this case the functions $e^{i\lambda\log_{t_{l1}}}$ extended to $b\mathbb{R}$ are identified with the extensions to $b\mathbb{R}$ of functions $c_{\lambda s}e^{i\lambda t}$ on $\mathbb{R}$, $c_{\lambda s} := e^{i\lambda\ln s}$, by the classical Bohr theorem there is $\lambda_0 \in \mathbb{R}$ such that the extension of $e^{i\lambda_0\log_{t_{l1}}}$ to $b\mathbb{R}$ separates $p_s(x)$ and $p_s(y)$. So, the extension of $e^{i\lambda_0\log_{t_{l1}}}$ to $\mathcal{M}(AP(\{z_l\}))$ separates x and y.

Suppose now that x and y are such that $p_s(x)$ and $p_s(y)$ belong to different Bohr compactifications of $\mathbb{R}$ for all s. This implies that x and y are limit points of the sets $\gamma_{t_l^{k(x)}}(s)$ and $\gamma_{t_l^{k(y)}}(s)$ for some $s \in (0, s_l)$ with $k(x) \neq k(y)$ and $k(x), k(y) \in \{-1, 1\}$. Then the function g_{t_l} by definition equals 1 at one of the points x, y and 0 at the other one.

Finally, assume that $x \in \mathcal{M}(AP(\{z_l\})) \setminus \partial\mathbb{D}$ and $y \in \partial\mathbb{D} \setminus \{z_l\}$. Then $g_{t_l}(x)$ equals either 0 or 1 and $g_{t_l}(y)$ differs from these numbers because g_{t_l} is decreasing on $\partial\mathbb{D} \setminus \{z_l\}$.

Thus we have proved that the family of functions $e^{i\lambda\log_{t_l^k}}$, g_{t_l} separate points on $\mathcal{M}(AP(\{z_l\}))$. This completes the proof of the theorem. $\square$

3.2. Proof of Corollary 1.5. According to Theorem 1.4 it suffices to prove that for functions g_{t_0} and $e^{i\lambda\log_{t_0^k}}$, $\lambda \in \mathbb{R}$, $z_0 := e^{it_0} \in \widetilde{S} := \phi(S)$, $k \in \{-1, 1\}$, the corresponding functions $\phi^*(g_{t_0})$ and $f := \phi^*(e^{i\lambda\log_{t_0^k}})$ belong to $AP(S)$. Since $g_{t_0} \in R_{\{z_0\}}$, the statement is trivial for $\phi^*(g_{t_0})$. Without loss of generality we may assume that ϕ preserves the orientation on $\partial\mathbb{D}$. Let $e^{i\tilde{t}_0} := \phi^{-1}(z_0)$. Then we have

$$\phi(e^{i(\tilde{t}_0+t)}) = e^{i(t_0+\tilde{\phi}(t))}, \quad t \in [0, 2\pi],$$

where $\tilde{\phi} : [0, 2\pi] \to [0, 2\pi]$ is a C^1 diffeomorphism and $\tilde{\phi}(0) = 0$. We set $c := \tilde{\phi}'(0)$. By the definition on the open arcs $\gamma_{\tilde{t}_0^k}(s)$ for $t \in (-\infty, 0)$ we obtain

$$\widehat{f}(t) := f(e^{i(\tilde{t}_0+kse^t)}) = e^{i\lambda\ln\tilde{\phi}(se^t))} = e^{i\lambda\ln(cse^t+o(se^t))} = e^{i(\lambda\ln(cs)+o(1))}e^{i\lambda t}, \quad \text{as} \quad s \to 0.$$

Since f is continuous outside $e^{i\tilde{t}_0}$, the latter implies that $f \in AP(S)$. $\square$

3.3. Proof of Theorem 1.6. We begin with the following

Lemma 3.1 *The algebra R_S is the uniform closure of the algebra generated by the algebras $R_{\{z\}}$ for all possible $z \in S$.*

(For $S = \partial\mathbb{D}$ a similar statement first was proved by Dieudonne [D]).

Proof. Consider a regulated function $f \in R_S$. Since $\partial\mathbb{D}$ is a compact set, by the definition of R_S for any $\epsilon > 0$ there are finitely many points $z_l := e^{it_l} \in \partial\mathbb{D}$, numbers

$s_l \in (0,\pi)$ and constant functions $f_k^l : \gamma_{t_l^k}(s_l) \to \mathbb{C}$, $k \in \{-1,1\}$, $1 \le l \le n$, such that

$$\bigcup_{l=1}^{n} (\gamma_{t_l^{-1}}(s_l) \cup \gamma_{t_l^1}(s_l)) = \partial\mathbb{D} \setminus \{z_1, \dots, z_n\} \quad \text{and for all} \quad 1 \le l \le n$$

$$\operatorname*{ess\,sup}_{z \in \gamma_{t_l^1}(s_l)} |f(z) - f_1^l(z)| < \epsilon \ , \qquad \operatorname*{ess\,sup}_{z \in \gamma_{t_l^{-1}}(s_l)} |f(z) - f_{-1}^l(z)| < \epsilon \ .$$

We set $U_l := \gamma_{t_l^1}(s_l) \cup \gamma_{t_l^{-1}}(s_l) \cup \{z_l\}$. Then $U = (U_l)_{l=1}^n$ is a finite open cover of $\partial\mathbb{D}$. Let $\{\rho_l\}_{l=1}^n$ be a continuous partition of unity subordinate to U such that $\operatorname{supp} \rho_l \subset\subset U_l$ and $\rho_l(z_l) = 1$, $1 \le l \le n$. Consider the functions f_l on $\partial\mathbb{D} \setminus \{z_l\}$ defined by the formulas

$$f_l(z) := \begin{cases} \rho_l(z) f_{-1}^l(z), & \text{if} \quad z \in \gamma_{t_l^{-1}}(s_l), \\ \\ \rho_l(z) f_1^l(z), & \text{if} \quad z \in \gamma_{t_l^1}(s_l). \end{cases}$$

If $z_l \in S$, then by definition $f_l \in R_{\{z_l\}}$. If $z_l \notin S$, then since f is continuous outside the compact set S, we can choose the above cover U and the family of functions $\{f_k^m\}_{1 \le m \le n}$, $k \in \{-1,1\}$, such that in the U_l the functions f_k^l, $k \in \{-1,1\}$, have the same limit at z_l. This implies the continuity of f_l on $\partial\mathbb{D}$, i.e., $f_l \in R_{\{z_l\}}$, as well. Also, we have

$$||f - \sum_{l=1}^{n} f_l||_{L^\infty(\partial\mathbb{D})} < \epsilon.$$

This completes the proof of the lemma. □

Let $F_1 \subset F_2$ be finite subsets of S. Then we have the natural injection $i_{F_1F_2} : R_{F_1} \hookrightarrow R_{F_2}$. Passing to the map dual to $i_{F_1F_2}$ we obtain a continuous map of the corresponding maximal ideal spaces: $p_{F_1F_2} : \mathcal{M}(R_{F_2}) \to \mathcal{M}(R_{F_1})$. Now the family $\{(\mathcal{M}(R_{F_1}), \mathcal{M}(R_{F_2}), p_{F_1F_2})\}_{F_1 \subset F_2 \subset S}$ determines an inverse limit system whose limit according to Lemma 3.1 coincides with $\mathcal{M}(R_S)$. By $p_F : \mathcal{M}(R_S) \to \mathcal{M}(R_F)$, $F \subset S$ is finite, we denote the limit maps determined by this limit. Then we have

$$c_F \circ p_F = c_S. \tag{3.3}$$

Let F consist of n points. Then $\partial\mathbb{D} \setminus F$ is a disjoint union of open arcs γ_k, $1 \le k \le n$. Consider a real function g_F on $\partial\mathbb{D}$ having discontinuities of the first kind at points F and continuous outside F such that the $g_F : \partial\mathbb{D} \setminus F \to \mathbb{R}$ is an injection and $g_F(\partial\mathbb{D} \setminus F)$ is the union of open intervals whose closures are mutually disjoint. Then an argument similar to that used in the proof of Theorem 1.4 (see the case $z_l \notin S$ there) shows that $R_F = \mathbb{C}(g_F)$, the uniform algebra in $L^\infty(\partial\mathbb{D})$ generated by g_F. This implies that $\mathcal{M}(R_F)$ is naturally homeomorphic to the disjoint union of closures $\overline{\gamma}_k$ of γ_k, $1 \le k \le n$, and $c_F : \mathcal{M}(R_F) \to \partial\mathbb{D}$ maps identically every γ_k in this union to $\gamma_k \subset \partial\mathbb{D}$. Since c_F is continuous, for every $z \in F$ the preimage $c_F^{-1}(z)$ consists of two points z_+ and z_- that can be naturally identified with counterclockwise and clockwise orientations of $\partial\mathbb{D}$ at z. Thus we obtain proofs of statement (1)-(3) of the theorem for a finite subset $F \subset S$. To prove the general case we use (3.3).

Assume that for some $z \in \partial\mathbb{D}$ the preimage $c_S^{-1}(z)$ contains at least three points x_1, x_2 and x_3. Then by the definition of the inverse limit, there is a finite subset $F \subset S$ such that $p_F(x_1)$, $p_F(x_2)$ and $p_F(x_3)$ are distinct points in $\mathcal{M}(R_F)$. Since by (3.3) the images of these points under c_F coincide with z, from the case established above for $\mathcal{M}(R_F)$ we obtain that $c_F^{-1}(z)$ consists of at most two points, a contradiction. Thus $c_S^{-1}(z)$ consists of at most 2 points for every $z \in \partial\mathbb{D}$.

Assume now that $z \in S$. Let $F \subset S$ be a finite subset containing z. Then the preimage $c_F^{-1}(z)$ consists of two points. Since by (3.3) $c_S^{-1}(z) = p_F^{-1}(c_F^{-1}(z))$, the preimage $c_S^{-1}(z)$ consists of two points, as well. (As before, they can be naturally identified with counterclockwise and clockwise orientations of $\partial\mathbb{D}$ at z.)

This proves statement (1) of the theorem.

Next, for $z \notin S$ we have $c_S^{-1}(z)$ is a single point. (For otherwise, for some finite F we will have that $c_F^{-1}(z)$ consists of at least 2 points, a contradiction.) Since $c_S^{-1}(S)$ is compact, the latter implies that $c_S : \mathcal{M}(R_S) \setminus c_S^{-1}(S) \to \partial\mathbb{D} \setminus S$ is a homeomorphism.

The proof of statement (2) is complete.

To prove (3) we will assume that S is infinite (for finite S the statement is already proved). Let $F \subset S$ be a finite subset consisting of n points, $n \geq 2$. Let $\partial\mathbb{D} \setminus F$ be the disjoint union of open arcs γ_k, $1 \leq k \leq n$. By χ_{γ_k} we denote the characteristic function of γ_k. Then every χ_{γ_k} belongs to R_F. By the same symbols we denote continuous extensions of χ_{γ_k} to $\mathcal{M}(R_F)$ by means of the Gelfand transform. Let us determine a continuous map $K_F : \mathcal{M}(R_F) \to \mathbb{Z}_2(F) := \{0,1\}^n$ by the formula

$$K_F(m) := (\chi_{\gamma_1}(m), \dots, \chi_{\gamma_n}(m)), \quad m \in \mathcal{M}(R_F).$$

Set $\mathcal{Z}(S) := \prod_{F \subset S} \mathbb{Z}_2(F)$ and determine the map $\mathcal{K}_S : \mathcal{M}(R_S) \to \mathcal{Z}(S)$ by the formula

$$\mathcal{K}_S(m) := \{K_F(p_F(m))\}_{F \subset S}.$$

We equip $\mathcal{Z}(S)$ with the *Tychonoff topology.* Then $\mathcal{Z}(S)$ is a totally disconnected compact Hausdorff space and the map $\mathcal{K}_S$ is continuous.

Show that $\mathcal{K}_S|_{c_S^{-1}(S)} : c_S^{-1}(S) \to \mathcal{Z}(S)$ is an injection. Indeed, for distinct points x, y from $c_S^{-1}(S)$ there exists a finite subset $F \subset S$ consisting of at least two points such that $p_F(x) \neq p_F(y)$ and $p_F(x), p_F(y) \in c_F^{-1}(F)$. Then by the definition of the map K_F we have

$$K_F(p_F(x)) \neq K_F(p_F(y)).$$

This implies that $\mathcal{K}_S(x) \neq \mathcal{K}_S(y)$. Since $c_S^{-1}(S)$ is a compact set, the injectivity implies that $c_S^{-1}(S)$ is homeomorphic to $\mathcal{K}_S(c_S^{-1}(S))$. The latter space is totally disconnected as a compact subset of the totally disconnected space $\mathcal{Z}(S)$.

This completes the proof of (3).

(4) According to (3.2) the maximal ideal space $\mathcal{M}(AP(S))$ of the algebra $AP(S)$ is homeomorphic to the inverse limit of compact spaces $\mathcal{M}(AP(F))$ with $F \subset S$ finite. Let $\widetilde{p}_{F_1F_2} : \mathcal{M}(AP(F_2)) \to \mathcal{M}(AP(F_1))$, $F_1 \subset F_2$, be continuous maps

determining this limit, and $\widetilde{p}_F : \mathcal{M}(AP(S)) \to \mathcal{M}(AP(F))$ be the corresponding limit maps. Since each $AP(F)$ is a self-adjoint algebra, by the Stone-Weierstrass theorem $\partial\mathbb{D} \setminus F$ is dense in $\mathcal{M}(AP(F))$. Hence, $\widetilde{p}_{F_1F_2}$ and $\widetilde{p}_F$ are surjective maps.

We begin with the description of $\mathcal{M}(AP(F))$. Suppose that $F := \{z_1, \dots, z_n\}$ and $F_i := F \setminus \{z_i\}$, $1 \le i \le n$. Consider disjoint union

$$X = \bigsqcup_{1\le i\le n} (\mathcal{M}(AP(\{z_i\})) \setminus F_i).$$

Note that each component of X contains $\partial\mathbb{D} \setminus F$ as an open subset. By $h_i : \partial\mathbb{D}\setminus F \hookrightarrow \mathcal{M}(AP(\{z_i\}))\setminus F_i$ we denote the corresponding embeddings. Then for each $z \in \partial\mathbb{D} \setminus F$ we sew together points $h_i(z)$, $1 \le i \le n$, and identify the obtained point with z. As a result we obtain the quotient space $\widetilde{X}$ of X and the "sewing“ map $\pi : X \to \widetilde{X}$. We equip $\widetilde{X}$ with the quotient topology:

$$U \subset \widetilde{X} \quad \text{is open} \qquad \Longleftrightarrow \qquad \pi^{-1}(U) \subset X \quad \text{is open.}$$

Lemma 3.2 *$\widetilde{X}$ is homeomorphic to $\mathcal{M}(AP(F))$.*

Proof. By definition each $V_i := \pi(\mathcal{M}(AP(\{z_i\})) \setminus F_i)$ is an open subset of $\widetilde{X}$ homeomorphic to $\mathcal{M}(AP(\{z_i\})) \setminus F_i$. Since the latter spaces are Hausdorff, $\widetilde{X}$ is Hausdorff, as well. Let us cover $\partial\mathbb{D}$ by closed arcs $\gamma_1, \dots, \gamma_n$ such that $\gamma_i \cap F = \{z_i\}$, $1 \le i \le n$. By $\widetilde{\gamma}_i$ we denote the closure of γ_i in $\mathcal{M}(AP(\{z_i\}))$. Then $\widetilde{\gamma}_i$ is a compact subset of $\mathcal{M}(AP(\{z_i\})) \setminus F_i$ and $U_i := \pi(\widetilde{\gamma}_i)$ is a compact subset of V_i. It is easy to see that $\widetilde{X} = \cup_{1\le i\le n} U_i$. Thus $\widetilde{X}$ is a compact space. Further, according to (3.2) each function from $AP(F)$ is extended continuously to $\widetilde{X}$ and the algebra of the extended functions separates points on $\widetilde{X}$. Hence by the Stone-Weierstrass theorem, $\widetilde{X}$ is homeomorphic to $\mathcal{M}(AP(F))$. $\square$

As a corollary of the above construction we immediately obtain the following:

Let $F_1 \subset F_2$ be finite subsets of S. Consider the commutative diagram

$$\begin{tikzcd}
\mathcal{M}(AP(S)) \arrow[r, "r_S"] \arrow[d, "\widetilde{p}_{F_2}"'] & \mathcal{M}(R_S) \arrow[r, "c_S"] \arrow[d, "p_{F_2}"'] & \partial\mathbb{D} \arrow[d, equal] \\
\mathcal{M}(AP(F_2)) \arrow[r, "r_{F_2}"] \arrow[d, "\widetilde{p}_{F_1F_2}"'] & \mathcal{M}(R_{F_2}) \arrow[r, "c_{F_2}"] \arrow[d, "p_{F_1F_2}"'] & \partial\mathbb{D} \arrow[d, equal] \\
\mathcal{M}(AP(F_1)) \arrow[r, "r_{F_1}"] & \mathcal{M}(R_{F_1}) \arrow[r, "c_{F_1}"] & \partial\mathbb{D}.
\end{tikzcd} \tag{3.4}$$

Here $\widetilde{p}_{F_1} := \widetilde{p}_{F_1F_2} \circ \widetilde{p}_{F_2}$ and $p_{F_1} := p_{F_1F_2} \circ p_{F_2}$. We set $\widetilde{F}_1 := (c_{F_1} \circ r_{F_1})^{-1}(F_1)$ and $\widetilde{S}_i := (c_{F_i} \circ r_{F_i})^{-1}(S)$, $i = 1, 2$. Then

(A)

$$\widetilde{p}_{F_1F_2} : \widetilde{p}^{\,-1}_{F_1F_2}(\widetilde{F}_1) \to \widetilde{F}_1$$

is a homeomorphism;

(B)

$$\mathcal{M}(AP(F_2)) \setminus \widetilde{S}_2 \xrightarrow{\widetilde{p}_{F_1F_2}} \mathcal{M}(AP(F_1)) \setminus \widetilde{S}_1 \xrightarrow{c_{F_1}\circ r_{F_1}} \partial\mathbb{D} \setminus S$$

are the identity maps.

From here by the definition of the inverse limit we obtain

(A1)

$$\widetilde{p}_{F_1} : \widetilde{p}_{F_1}^{-1}(\widetilde{F}_1) \to \widetilde{F}_1$$

is a homeomorphism;

(B1)

$$\mathcal{M}(AP(S)) \setminus (c_S \circ r_S)^{-1}(S) \xrightarrow{\widetilde{p}_{F_1}} \mathcal{M}(AP(F_1)) \setminus \widetilde{S}_1 \xrightarrow{c_{F_1}\circ r_{F_1}} \partial\mathbb{D} \setminus S$$

are the identity maps.

Assume now that $F_1 = \{z\} \subset S$. Then according to Theorem 1.4 (1) the set $c_{F_1}^{-1}(F_1) = c_S^{-1}(F_1)$ consists of two points $\{z_+\}$ and $\{z_-\}$ identified with counterclockwise and clockwise orientations at z. Thus to prove (4) we must show (according to (3.4) and statements (A), (A1)) that each set $r_{\{z\}}^{-1}(z_\pm)$ is homeomorphic to $b\mathbb{R}$.

To this end let us recall that in the proof of Theorem 1.4 we had established that $\mathcal{M}(AP(\{z\}))$ is the inverse limit of the maximal ideal spaces $\mathcal{M}_{\{z\}}(s)$ of algebras $AP_{\{z\}}(s)$ of continuous functions on $\partial\mathbb{D}\setminus\{z\}$ almost periodic on the open arcs $\gamma_{t^k}(s)$ where $z := e^{it}$ and $s \in (0,\pi)$. Also, in that proof the structure of each $\mathcal{M}_{\{z\}}(s)$ was described.

For every pair $0 < s'' < s' < \pi$ let $p_{s''s'} : \mathcal{M}_{\{z\}}(s'') \to \mathcal{M}_{\{z\}}(s')$ be the continuous surjective map dual to the embedding $i_{s''s'} : AP_{\{z\}}(s') \hookrightarrow AP_{\{z\}}(s'')$. From the proof of Theorem 1.4 we know that every $\mathcal{M}_{\{z\}}(s)$ is obtained by gluing $\partial\mathbb{D} \setminus \{z\}$ with two copies of $b\mathbb{R}$ where one copy (denoted $b\mathbb{R}_1$) is obtained by gluing with $\gamma_{t^1}(s)$ and another one (denoted by $b\mathbb{R}_{-1}$) is obtained by gluing with $\gamma_{t^{-1}}(s)$. Suppose that $\xi \in b\mathbb{R}_1 \subset \mathcal{M}_{\{z\}}(s'')$. Let us compute $p_{s''s'}(\xi) \in b\mathbb{R}_1 \subset \mathcal{M}_{\{z\}}(s')$. Let $\{z_\alpha\} \subset \gamma_{t^1}(s'')$ be a net converging to ξ. This means that the net $\{\phi_{s''}(z_\alpha)\} \subset \mathbb{R}_-$ converges to ξ in the topology of the Bohr compactification on $b\mathbb{R}$; here $\phi_{s''}$ is the map inverse to the map $\psi_{s''} : \mathbb{R}_- \to \gamma_{t^1}(s'')$, $x \mapsto e^{i(t+s''e^x)}$. Next, by the definition the net $\{\phi_{s'}(z_\alpha)\}$ converges to $p_{s''s'}(\xi)$. A straightforward computation shows that

$$\phi_{s'}(z_\alpha) = \phi_{s''}(z_\alpha) + \ln\left(\frac{s'}{s''}\right) \quad \text{for all} \quad z_\alpha.$$

Thus we have

$$p_{s''s'}(\xi) = \xi + \ln\left(\frac{s'}{s''}\right), \quad \xi \in b\mathbb{R}_1. \tag{3.5}$$

Here the sum denotes the group operation on $b\mathbb{R}$. Similarly,

$$p_{s''s'}(\xi) = \xi + \ln\left(\frac{s'}{s''}\right), \quad \xi \in b\mathbb{R}_{-1}. \tag{3.6}$$

Using these formulas we now prove that each $r^{-1}_{\{z\}}(z_\pm)$ is homeomorphic to $b\mathbb{R}$. We will prove the statement for z_+ (for z_- the argument is similar).

For a fixed $s_0 \in (0,\pi)$ consider the limit map $p_{s_0} : \mathcal{M}(AP(\{z\})) \to \mathcal{M}_{\{z\}}(s_0)$. Then p_{s_0} maps $r^{-1}_{\{z\}}(z_+)$ into $X_{s_0} := b\mathbb{R}_1 \subset \mathcal{M}_{\{z\}}(s_0)$. Moreover, by definition $r^{-1}_{\{z\}}(z_+)$ is the inverse limit of the system $\{(X_{s''}, X_{s'}, p_{s''s'})\}$ where we write X_s for $b\mathbb{R}_1 \subset \mathcal{M}_{\{z\}}(s)$. Since according to (3.5) every $p_{s''s'} : X_{s''} \to X_{s'}$ is a homeomorphism (even an automorphism of $b\mathbb{R}$), by the definition of the inverse limit $p_{s_0} : r^{-1}_{\{z\}}(z_+) \to X_{s_0}$ is a homeomorphism.

This completes the proof of statement (4).

(5) The required statement follows from (B1) and Theorem 1.6 (2).

The proof of Theorem 1.6 is complete. $\square$

Remark 3.3 It is well known that the covering dimension of $b\mathbb{R}$ is ∞ because this group is the inverse limit of compact abelian Lie groups whose dimensions tend to ∞. Since $b\mathbb{R} \subset \mathcal{M}(AP(S))$, the covering dimension of $\mathcal{M}(AP(S))$ is ∞, as well (statement (6)). Also, statement (7) follows from the fact that $c_S^{-1}(S)$ is totally disconnected. Indeed, for a continuous map $\phi : T \to (c_S \circ r_S)^{-1}(S)$ of a connected topological space T the image of $r_S \circ \phi$ is a single point. This implies the required.

4. Proofs of Theorem 1.7 and Example 1.9

4.1. In this part we formulate and prove some auxiliary results used in the proof of the theorem.

Notation. *Let $z_0 \in \partial\mathbb{D}$ and U_{z_0} be the intersection of an open disk of radius ≤ 1 centered at z_0 with $\overline{\mathbb{D}} \setminus \{z_0\}$. We call such U_{z_0} a circular neighbourhood of z_0.*

Next, we define almost periodic continuous functions on a circular neighbourhood U_{z_0} of z_0 holomorphic in its interior points as follows.

Let $\phi_{z_0} : \mathbb{D} \to \mathbb{H}_+$,

$$\phi_{z_0}(z) := \frac{2i(z_0 - z)}{z_0 + z}, \quad z \in \mathbb{D}, \tag{4.1}$$

be a conformal map of $\mathbb{D}$ onto the upper half-plane $\mathbb{H}_+$. Then ϕ_{z_0} is also continuous on $\partial\mathbb{D} \setminus \{-z_0\}$ and maps it diffeomorphically onto $\mathbb{R}$ (the boundary of $\mathbb{H}_+$) so that $\phi_{z_0}(z_0) = 0$. Let Σ_0 be the interior of the strip $\Sigma := \{z \in \mathbb{C} \; : \; \text{Im } z \in [0,\pi]\}$. Consider the conformal map $\text{Log} : \mathbb{H}_+ \to \Sigma_0$, $z \mapsto \text{Log}(z) := \ln|z| + i\text{Arg}(z)$, where $\text{Arg} : \mathbb{C} \setminus \mathbb{R}_- \to (-\pi,\pi)$ is the principal branch of the multi-function arg, the argument of a complex number. The function Log is extended to a homeomorphism of $\overline{\mathbb{H}}_+ \setminus \{0\}$ onto Σ; here $\overline{\mathbb{H}}_+$ stands for the closure of $\mathbb{H}_+$.

By $AP_{\mathcal{C}}(\Sigma)$ we denote the algebra of uniformly continuous almost periodic functions on Σ (i.e., they are almost periodic on any line parallel to the x-axis). Clearly we have $AP_{\mathcal{O}}(\Sigma) \subset AP_{\mathcal{C}}(\Sigma)$. Then according to Theorem 2.4 (the corona theorem for $AP_{\mathcal{O}}(\Sigma)$), the maximal ideal space $\mathcal{M}(AP_{\mathcal{C}}(\Sigma))$ of the algebra $AP_{\mathcal{C}}(\Sigma)$ is homeomorphic to $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$. In what follows we identify these spaces.

Definition 4.1 *We say that* $f : U_{z_0} \to \mathbb{C}$ *is a (continuous) almost periodic function if there is a function* $\widehat{f} \in AP_{\mathcal{C}}(\Sigma)$ *such that*

$$f(z) := \widehat{f}(\mathrm{Log}(\phi_{z_0}(z))) \quad \text{for all} \quad z \in U_{z_0}.$$

If such $\widehat{f} \in AP_{\mathcal{O}}(\Sigma)$, *then* f *is called a holomorphic almost periodic function.*

Suppose that $z_0 = e^{it_0}$. For $s \in (0, \pi)$ we set $\gamma_1(z_0, s) := \mathrm{Log}(\phi_{z_0}(\gamma_{t_0^1}(s))) \subset \mathbb{R}$ and $\gamma_{-1}(z_0, s) := \mathrm{Log}(\phi_{z_0}(\gamma_{t_0^{-1}}(s))) \subset \mathbb{R} + i\pi$.

Lemma 4.2 *Let* $f \in AP(\{-z_0, z_0\})$. *We set* $f_k := f|_{\gamma_{t_0^k}(\pi)}$ *and consider the functions* $h_k = f_k \circ \varphi_{z_0}^{-1} \circ \mathrm{Log}^{-1}$ *on* $\gamma_k(z_0, s)$, $k \in \{-1, 1\}$. *Then for any* $\epsilon > 0$ *there are* $s_\epsilon \in (0, s)$ *and almost periodic functions* h'_1 *on* $\mathbb{R}$, *and* h'_{-1} *on* $\mathbb{R} + i\pi$ *such that*

$$\sup_{z \in \gamma_k(z_0, s_\epsilon)} |h_k(z) - h'_k(z)| < \epsilon \quad \text{for each} \quad k \in \{-1, 1\}. \tag{4.2}$$

Proof. We prove the result for f_1 only. The proof for f_{-1} is similar. According to Theorem 1.4 it suffices to prove the lemma for $f_1 = g_{t_0}$ or $f_1 = e^{i\lambda \log_{t_0^1}}$, $\lambda \in \mathbb{R}$. In the first case we can choose a sufficiently small s_ϵ such that on $\gamma_{t_0^1}(s_\epsilon)$ the function g_{t_0} is uniformly approximated with an error $< \epsilon$ by a constant function. Then as h'_1 we can choose the corresponding constant function on $\gamma_1(z_0, s_\epsilon)$. In the second case, by definition,

$$h_1(x) = e^{i\lambda \ln\left(\mathrm{Arg}\left(\frac{2i - e^x}{2i + e^x}\right)\right)} = e^{i\lambda \ln\left(\frac{4e^x}{4 - e^{2x}} + o(e^{3x})\right)} = e^{i\lambda(x + o(e^x))} \quad \text{as} \quad x \to -\infty.$$

From here for a sufficiently small s_ϵ we have

$$|h_1(x) - e^{i\lambda x}| < \epsilon \quad \text{for all} \quad x \in \gamma_1(z_0, s_\epsilon). \qquad \square$$

We also use the following well-known result.

Lemma 4.3 *Suppose that* f_1 *and* f_2 *are continuous almost periodic functions on* $\mathbb{R}$ *and* $\mathbb{R} + i\pi$, *respectively. Then there exists a function* $F \in AP_{\mathcal{C}}(\Sigma)$ *harmonic in* Σ_0 *whose boundary values are* f_1 *and* f_2.

Proof. Let F be a function harmonic in Σ_0 with boundary values f_1 and f_2. Since f_1 and f_2 are almost periodic, for any $\epsilon > 0$ there exists $l(\epsilon) > 0$ such that every interval $[t_0, t_0 + l(\epsilon)]$ contains a common ϵ-period of f_1 and f_2, say, τ_ϵ, see, e.g., [LZ]. Thus

$$\sup_{x \in \mathbb{R}} |f_1(x + \tau_\epsilon) - f_1(x)| < \epsilon \quad \text{and} \quad \sup_{x \in \mathbb{R}} |f_2(x + i\pi + \tau_\epsilon) - f_2(x + i\pi)| < \epsilon.$$

Now, by the maximum principle for harmonic functions

$$\sup_{x \in \mathbb{R}} |F(x + iy + \tau_\epsilon) - F(x + iy)| < \epsilon \quad \text{for each} \quad y \in [0, \pi],$$

that is, F is almost periodic on every line $\mathbb{R} + iy$, $y \in [0, \pi]$. $\square$

Let $\mathcal{A}(U_{z_0})$ be the algebra of continuous functions on $\mathbb{D}$ almost periodic on the circular neighbourhood U_{z_0} of z_0. By $\mathcal{A}_{z_0}$ we denote the uniform closure of the algebra generated by all $\mathcal{A}(U_{z_0})$ and by $\mathcal{M}_{z_0}$ the closure of $\mathbb{D}$ in the maximal ideal space of $\mathcal{A}_{z_0}$. Since the algebra $\mathcal{A}_{z_0}$ is self-adjoint, by the Stone-Weierstrass theorem $\mathcal{M}_{z_0}$ coincides with the maximal ideal space of $\mathcal{A}_{z_0}$. Next, let $p_{z_0} : \mathcal{M}_{z_0} \to \overline{\mathbb{D}}$ be the continuous surjective map dual to the natural embedding $C(\overline{\mathbb{D}}) \hookrightarrow \mathcal{A}_{z_0}$. Then we have

Lemma 4.4 *(a) For every neighbourhood U of the compact set $F_{z_0} := p_{z_0}^{-1}(z_0)$ there is a circular neighbourhood U_{z_0} of z_0 such that $U_{z_0} \cap \mathbb{D} \subset U \cap \mathbb{D}$.*

(b) F_{z_0} is homeomorphic to $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$.

(c) Each function $f \in AP(S) \cap H^\infty$ belongs to the algebra $\bigcap_{z \in \partial\mathbb{D}} \mathcal{A}_z$.

Proof. (a), (b). Since the algebra $\mathcal{A}(U_{z_0})$ is self-adjoint, $\mathbb{D}$ is dense in its maximal ideal space $\mathcal{M}(U_{z_0})$. Then $\mathcal{M}_{z_0}$ is the inverse limit of the compact spaces $\mathcal{M}(U_{z_0})$ (because $\mathcal{A}_{z_0}$ is the uniform closure of the algebra generated by algebras $\mathcal{A}(U_{z_0})$), see, e.g, [R]. For $U_{z_0} \subset V_{z_0}$ by $p_{U_{z_0}V_{z_0}} : \mathcal{M}(U_{z_0}) \to \mathcal{M}(V_{z_0})$ we denote the maps in this limit system and by $p_{U_{z_0}} : \mathcal{M}_{z_0} \to \mathcal{M}(U_{z_0})$ the corresponding (continuous and surjective) limit maps. Then, by the definition of the inverse limit, the base of topology on $\mathcal{M}_{z_0}$ consists of the sets $p_{U_{z_0}}^{-1}(U)$ where $U \subset \mathcal{M}(U_{z_0})$ is open and U_{z_0} is a circular neighbourhood of z_0. In particular, since F_{z_0} is a compact set, for a neighbourhood U of F_{z_0} there is a circular neighbourhood $\widetilde{U}_{z_0}$ of z_0 and a neighbourhood $\widetilde{U} \subset \mathcal{M}(\widetilde{U}_{z_0})$ of $F(\widetilde{U}_{z_0}) := p_{\widetilde{U}_{z_0}}(F_{z_0})$ such that $p_{\widetilde{U}_{z_0}}^{-1}(\widetilde{U}) \subset U$. Recall that $\mathbb{D}$ is a dense subset of $\mathcal{M}(\widetilde{U}_{z_0})$ and $\mathcal{M}_{z_0}$. Also, $p_{\widetilde{U}_{z_0}}^{-1}(\mathbb{D})$ contains $\mathbb{D} \subset \mathcal{M}_{z_0}$. Thus in order to prove (a) it suffices to show that there is $U_{z_0} \subset \widetilde{U}_{z_0}$ such that $U_{z_0} \cap \mathbb{D} \subset \widetilde{U} \cap \mathbb{D}$.

First, let us study the structure of $\mathcal{M}(\widetilde{U}_{z_0})$. Let $\mathcal{A}^*(\widetilde{U}_{z_0})$ be the pullback to Σ by means of the map $(\mathrm{Log} \circ \phi_{z_0})^{-1}$ of the algebra $\mathcal{A}(\widetilde{U}_{z_0})$. Then $\mathcal{A}^*(\widetilde{U}_{z_0})$ consists of continuous functions on Σ_0 such that on $(\mathrm{Log} \circ \phi_{z_0})(\widetilde{U}_{z_0})$ they are restrictions of almost periodic functions on Σ. Since $\mathcal{A}^*(\widetilde{U}_{z_0})$ is isomorphic to $\mathcal{A}(\widetilde{U}_{z_0})$ we can naturally identify the maximal ideal spaces of these algebras.

Further, observe that there is $T < 0$ such that $(\mathrm{Log} \circ \phi_{z_0})(\widetilde{U}_{z_0})$ contains the subset $\Sigma_T := \{z \in \Sigma \ : \ \mathrm{Re}\ z \le T\}$ of the strip Σ. By the definition of the topology on $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ (see section 2) Σ_T is dense in $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$. Hence, the space $\mathcal{M}(\widetilde{U}_{z_0})$ contains $\mathcal{M}(AP_{\mathcal{C}}(\Sigma))(= \mathcal{M}(AP_{\mathcal{O}}(\Sigma)))$. Let K be the intersection of the closures of $\widetilde{U}_{z_0}$ and $\mathbb{D} \setminus \widetilde{U}_{z_0}$ in $\mathbb{C}$. Then $K' := \mathrm{Log} \circ \phi_{z_0}(K)$ is a compact subset of Σ. In particular, $AP_{\mathcal{C}}(\Sigma)|_{K'} = C(K')$. This implies (by the Tietze extension theorem) that every bounded continuous function on $(\mathbb{D} \setminus \widetilde{U}_{z_0}) \cup K$ can be extended to a function from $\mathcal{A}(\widetilde{U}_{z_0})$ with the same supremum norm. Now, we can describe explicitly $\mathcal{M}(\widetilde{U}_{z_0})$ as follows (cf. the proof of Theorem 1.4 for a similar construction).

Let M be the maximal ideal space of the algebra of bounded continuous functions on $(\mathbb{D} \setminus \widetilde{U}_{z_0}) \cup K$. We identify $K \subset M$ with $K' \subset \mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ by means of

$\mathrm{Log}\circ\phi_{z_0}$. Then the quotient space of $M\sqcup\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ under this identification is homeomorphic to $\mathcal{M}(\widetilde{U}_{z_0})$.

Next, by definition the fibre F_{z_0} over z_0 consists of limit points in $\mathcal{M}_{z_0}$ of all nets converging to $\{z_0\}$ inside $\mathbb{D}$. This and the above construction of $\mathcal{M}(\widetilde{U}_{z_0})$ show that $F(\widetilde{U}_{z_0})$ is homeomorphic to $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$. Moreover, F_{z_0} is the inverse limit of compact sets $F(\widetilde{U}_{z_0})$ where the limit system is determined by the maps $p_{U_{z_0}V_{z_0}}|_{F(U_{z_0})}$ for $U_{z_0}\subset V_{z_0}$. Let us show that the maps $p_{U_{z_0}V_{z_0}}|_{F(U_{z_0})}:F(U_{z_0})\to F(V_{z_0})$ are homeomorphisms. Indeed, let $\{z_\alpha\}\subset U_{z_0}$ be a net converging to a point $\xi\in F(U_{z_0})$. Since $U_{z_0}\hookrightarrow V_{z_0}$, in our definitions of $\mathcal{M}(U_{z_0})$ and $\mathcal{M}(V_{z_0})$ the net $\{z_\alpha\}$ converges to the same point $\xi\in F(V_{z_0})$ which gives the required statement. Since all maps $p_{U_{z_0}V_{z_0}}|_{F(U_{z_0})}$ are homeomorphisms, by the definition of the inverse limit the map $p_{\widetilde{U}_{z_0}}|_{F_{z_0}}:F_{z_0}\to F(\widetilde{U}_{z_0})$, is also a homeomorphism. This completes the proof of (b).

Now, observe that in our model of $\mathcal{M}(\widetilde{U}_{z_0})$ the intersection of $\widetilde{U}$ with $\mathbb{D}$ contains $\widetilde{U}_{z_0}\cap\mathbb{D}$, this completes the proof of (a).

(c) Fix a point $z_*\in\partial\mathbb{D}$. We must show that every $f\in AP(S)\cap H^\infty$ belongs to $\mathcal{A}_{z_*}$. According to (3.2) and Lemma 4.2 each $f\in AP(\partial\mathbb{D})\cap H^\infty$ can be approximated locally on open arcs of the form $\gamma_{t^k}(s)$, $k\in\{-1,1\}$, $s\in(0,\pi)$, $z:=e^{it}\in\partial\mathbb{D}$, by pullbacks of almost periodic functions on the boundary of Σ. Using compactness of $\partial\mathbb{D}$, for a given $\epsilon>0$ we can find a finitely many points $z_1,\dots,z_n\in\partial\mathbb{D}$, arcs $\gamma_{t_l^k}(s_l)$, $k\in\{-1,1\}$, $s_l\in(0,\pi)$, $z_l:=e^{it_l}$, and functions $f^l:\partial\mathbb{D}\setminus\{-z_l,z_l\}\to\mathbb{C}$ which are pullbacks of almost periodic functions on $\partial\Sigma$ by means of $\mathrm{Log}\circ\phi_{z_l}$, $1\le l\le n$, such that

$$\partial\mathbb{D}\setminus\{z_1,\dots,z_n\}=\bigcup_{1\le l\le n}(\gamma_{t_l^{-1}}\cup\gamma_{t_l^1})\quad\text{and for all}\quad 1\le l\le n$$

$$\operatorname*{ess\,sup}_{z\in\gamma_{t_l^1}(s_l)\cup\gamma_{t_l^{-1}}(s_l)}|f(z)-f^l(z)|<\epsilon.$$

Without loss of generality we may assume that $z_*\in\{z_1,\dots,z_n\}$. Next, set $V_l:=\gamma_{t_l^{-1}}\cup\gamma_{t_l^1}\cup\{z_l\}$. Then $(V_l)_{l=1}^n$ is a finite open cover of $\partial\mathbb{D}$. Let $\{\rho_l\}_{l=1}^n$ be a smooth partition of unity subordinate to this cover such that $\rho_l(z_l)=1$. Consider the functions f_l on $\partial\mathbb{D}\setminus\{z_l\}$ defined by the formulas

$$f_l:=\rho_l f^l,\quad 1\le l\le n.$$

Let F be the harmonic function on $\mathbb{D}$ such that $F|_{\partial\mathbb{D}}=\sum_{1\le l\le n}f_l$. Then

$$||f-F||_{L^\infty(\mathbb{D})}<\epsilon.$$

We prove that $F\in\mathcal{A}_{z_*}$. Since $\epsilon>0$ is arbitrary, this will complete the proof of (c).

By $F_{l,1}$ and $F_{l,2}$ we denote the harmonic functions on $\mathbb{D}$ with the boundary values f_l and f^l-f_l, respectively. Thus $F_l:=F_{l,1}+F_{l,2}$ is the harmonic function with the boundary values f^l. According to Lemma 4.3, every F_l is almost periodic on $\overline{\mathbb{D}}\setminus\{\pm z_l\}$. Thus if $z_l=z_*$, then $F_l\in\mathcal{A}_{z_*}$. If $z_l\ne z_*$, then F_l is continuous at z_* and so by the definition of $\mathcal{A}_{z_*}$ the function $F_l\in\mathcal{A}_{z_*}$, as well. Further, for a point z_l distinct from z_* the function $F_{l,1}$ can be extended continuously in an open disk centered at z_* (because the support of f_l does not contain z_*). Hence, such $F_{l,1} \in \mathcal{A}_{z_*}$. Assume now that $z_l = z_*$ for some l. Then the function $F_{l,2}$ can be extended continuously in an open disk centered at z_* (because the support of $f^l - f_l$ does not contain z_*). Thus $F_{l,2} \in \mathcal{A}_{z_*}$ and in this case $F_{l,1} := F_l - F_{l,2} \in \mathcal{A}_{z_*}$, as well. Since $F := \sum_{1\le l\le n} F_{l,1}$, combining the above considered cases we obtain that $F \in \mathcal{A}_{z_*}$.

This completes the proof of Lemma 4.4. $\square$

Theorem 4.5 *Let $f \in AP(S) \cap H^\infty$. Then for each $z_0 \in \partial\mathbb{D}$ and any $\epsilon > 0$ there is a circular neighbourhood $U_{z_0} := U_{z_0}(f, \epsilon)$ of z_0 and a holomorphic almost periodic function f_{z_0} on U_{z_0} such that*

$$\sup_{z\in U_{z_0}\cap\mathbb{D}} |f(z) - f_{z_0}(z)| < \epsilon.$$

Proof. Fix a point $z_0 \in \partial\mathbb{D}$. According to Lemma 4.4(c) the function $f \in AP(S) \cap H^\infty$ belongs to $\mathcal{A}_{z_0}$ and so it is extended by means of the Gelfand transform to a continuous function $\widehat{f}$ on $\mathcal{M}_{z_0}$. We use the description of $\mathcal{M}_{z_0}$ presented in the proof of Lemma 4.4. Recall that in that construction the fibre $F_{z_0} \subset \mathcal{M}_{z_0}$ over z_0 is naturally identified with $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$. Then we have (see Definition 2.1).

Lemma 4.6 *The function $\widehat{f}|_{F_{z_0}}$ is holomorphic.*

Proof. In the proof we use the results of section 2. Let us consider the map $i_\xi : \Sigma_0 \to \mathcal{M}(AP_{\mathcal{O}})(\Sigma)$, $\xi \in b\mathbb{Z}$. We must show that $\widehat{f} \circ i_\xi$ is holomorphic.

To this end we transfer the function f by means of the map $(\mathrm{Log} \circ \phi_{z_0})^{-1}$ to Σ_0 and by $\widetilde{f}$ denote the pulled back function. Fix a point $\eta \in i_\xi(\Sigma_0)$, say, $\eta := i_\xi(w)$, $w \in \Sigma_0$. Then there is a straight line $\mathbb{R}+iy \subset \Sigma_0$, $y \in (0, \pi)$, and a net $\{z_\alpha\} \subset \mathbb{R}+iy$ which forms an infinite discrete subset of $\mathbb{R}+iy$ such that $\{z_\alpha\}$ converges to η in the topology of $\mathcal{M}(AP_{\mathcal{O}})(\Sigma)$. Let $B \subset \mathbb{C}$ be an open disk such that $\{\mathbb{R}+iy\} + B \subset \Sigma_0$. By the definition of i_ξ, for each $z \in B$ the net $\{z_\alpha + z\}$ converges in $\mathcal{M}(AP_{\mathcal{O}})(\Sigma)$ to $i_\xi(w + z)$. Also, by the definition of $\mathcal{M}_{z_0}$ we have

$$\lim_\alpha \widetilde{f}(z_\alpha + z) = (\widehat{f} \circ i_\xi)(w + z), \quad z \in B.$$

But the holomorphic functions $\widetilde{f}_\alpha(z) := \widetilde{f}(z_\alpha + z)$ form a normal family on B. Therefore using an argument similar to that of the proof of Lemma 2.3 (1) we obtain that $\widehat{f} \circ i_\xi|_B$ is holomorphic. Since ξ and η are arbitrary, the latter implies that $\widehat{f}|_{F_{z_0}}$ is holomorphic. $\square$

Now by Lemma 2.3 we obtain that there is a function $\widetilde{f}_{z_0} \in AP_{\mathcal{O}}(\Sigma)$ whose extension to $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ coincides with $\widehat{f}|_{F_{z_0}}$. Let us consider the function $f_{z_0} \in \mathcal{A}_{z_0}$ whose pullback to Σ by means of $(\mathrm{Log} \circ \phi_{z_0})^{-1}$ coincides with $\widetilde{f}_{z_0}$. Then by the definition of the topology of $\mathcal{M}_{z_0}$ the extension $\widehat{f}_{z_0}$ of f_{z_0} to $\mathcal{M}_{z_0}$ satisfies $\widehat{f}_{z_0}|_{F_{z_0}} = \widehat{f}|_{F_{z_0}}$. Since F_{z_0} is a compact set, the latter implies that there is a neighbourhood U of F_{z_0}

in $\mathcal{M}_{z_0}$ such that $|\widehat{f}_{z_0}(x) - \widehat{f}(x)| < \epsilon$ for all $x \in U$. Finally, by Lemma 4.4(a) there is a circular neighbourhood U_{z_0} such that $U_{z_0} \cap \mathbb{D} \subset U \cap \mathbb{D}$. Thus $|f_{z_0}(z) - f(z)| < \epsilon$ for all $z \in U_{z_0} \cap \mathbb{D}$. $\Box$

4.2. Proof of Theorem 1.7. We must show that $A_S = AP(S) \cap H^\infty$. We split the proof into several parts. First we prove the following statement.

Lemma 4.7 *$AP(S) \cap H^\infty$ is the uniform closure of the algebra generated by all possible subalgebras $AP(F) \cap H^\infty$ with finite $F \subset S$.*

Then we will prove that $A_F = AP(F) \cap H^\infty$ for every finite subset $F \subset \partial\mathbb{D}$. Together with the above lemma and the fact that A_S is the uniform closure of the algebra generated by all possible subalgebras A_F with finite $F \subset S$, this will clearly complete the proof of the theorem.

Proof. According to Theorem 4.5, for a given $f \in AP(S) \cap H^\infty$ we can find finitely many points $z_1, \dots, z_n$, circular neighbourhoods $U_{z_1}, \dots, U_{z_n}$ and holomorphic almost periodic functions $f_1, \dots, f_n$ defined on $U_{z_1}, \dots, U_{z_n}$, respectively, such that $(U_{z_i})_{1 \le i \le n}$ forms an open cover of $\partial\mathbb{D} \setminus \{z_1, \dots, z_n\}$ and

$$\max_i ||f|_{U_{z_i}} - f_i||_{L^\infty(U_{z_i})} < \epsilon.$$

Since the discontinuities of $f|_{\partial\mathbb{D}}$ belong to the closed set S, each function f_i with $z_i \notin S$ can be chosen also to be continuous on the closure $\overline{U}_{z_i}$.

Further, form a cocycle $\{c_{ij}\}$ on the intersections of sets from the above cover by the formula

$$c_{ij}(z) := f_i(z) - f_j(z), \quad z \in U_{z_i} \cap U_{z_j}.$$

Diminishing, if necessary, the sets of the above cover we may assume without loss of generality that all $U_{z_i} \cap U_{z_j}$, $i \neq j$, do not contain points $z_1, \dots, z_n$. Then each $U_{z_i} \cap U_{z_j}$, $i \neq j$, is a compact subset of $\overline{\mathbb{D}}$ and the corresponding c_{ij} are continuous and holomorphic in interior points of $U_{z_i} \cap U_{z_j}$.

Let $\{\rho_i\}$ be a smooth partition of unity subordinate to the cover $(U_{z_i})_{1 \le i \le n}$. We can choose every ρ_i so that it is the restriction to $\overline{U}_{z_i}$ of a C^∞-function on $\mathbb{C}$ and $\rho_i(z_i) = 1$. As usual, we resolve the cocycle $\{c_{ij}\}$ using this partition of unity by the formulas

$$\widetilde{f}_j(z) = \sum_{k=1}^n \rho_k(z) c_{jk}(z), \quad z \in \overline{U}_{z_j}. \tag{4.3}$$

Hence,

$$c_{ij}(z) := \widetilde{f}_i(z) - \widetilde{f}_j(z), \quad z \in U_{z_i} \cap U_{z_j}.$$

In particular, since c_{ij} are holomorphic in $\mathbb{D} \cap U_{z_i} \cap U_{z_j}$, the formula

$$h(z) := \frac{\partial \widetilde{f}_i(z)}{\partial \overline{z}}, \quad z \in U_{z_i} \cap \mathbb{D},$$

determines a smooth bounded function in an open annulus $A \subset \cup_{i=1}^n \overline{U}_{z_i}$ with the outer boundary $\partial\mathbb{D}$. Also, by our choice of the partition of unity, h is extended continuously to the closure $\overline{A}$ of A.

Let us consider a function H determined by the formula

$$H(z) = \frac{1}{2\pi i} \int\int_{\zeta \in A} \frac{h(\zeta)}{\zeta - z}\, d\zeta \wedge d\overline{\zeta}, \quad z \in \overline{A}. \tag{4.4}$$

Passing in (4.4) to polar coordinates with the origin at z, we easily obtain

$$\sup_{z \in A} |H(z)| \le Cw(A) \sup_{z \in A} |h(z)| \tag{4.5}$$

where $w(A)$ is the width of A and $C > 0$ is an absolute constant. Moreover, $H \in C(\overline{A})$ and $\partial H/\partial \overline{z} = h$ in A, see, e.g., [G, Chapter VIII]. Let us replace A by a similar annulus of a smaller width such that for this new A

$$\sup_{z \in A} |H(z)| < \epsilon.$$

Now we set

$$c_i(z) := \widetilde{f}_i(z) - H(z), \quad z \in \overline{U}_{z_i} \cap \overline{A}.$$

Then each c_i is continuous on $U_{z_i} \cap \overline{A}$ holomorphic in interior points of this set and

$$c_i(z) - c_j(z) = c_{ij}(z), \quad z \in U_{z_i} \cap U_{z_j}.$$

Since every $|c_{ij}(z)| < 2\epsilon$ for all $z \in U_{z_i} \cap U_{z_j}$,

$$|c_i(z)| < 3\epsilon, \quad z \in U_{z_i} \cap U_{z_j}.$$

Let us determine a global function f_ϵ on $\overline{A} \setminus \{z_1, \dots, z_n\}$ by the formulas

$$f_\epsilon(z) := f_i(z) - c_i(z), \quad z \in U_i \cap \overline{A}.$$

Since for $z_i \notin S$, the function f_i is continuous on $\overline{U}_{z_i}$, from the above construction we obtain that $f_\epsilon \in H^\infty(A) \cap AP(F)$ where $F := \{z_1, \dots, z_n\} \cap S$. Also,

$$||f - f_\epsilon||_{L^\infty(A)} < 4\epsilon.$$

Let B be an open disk centered at 0 whose intersection with A is an annulus of width $< \epsilon$. Consider the cocycle c on $B \cap A$ defined by

$$c(z) = f(z) - f_\epsilon(z), \quad z \in B \cap A.$$

By definition, $|c(z)| \le 4\epsilon$ for all $z \in B \cap A$. Let A' be the open annulus with the interior boundary coinciding with the interior boundary of A and with the outer boundary $\{z \in \mathbb{C} \ : \ |z| = 2\}$. Then $A' \cap B = A \cap B$. Consider a smooth partition of unity subordinate to the cover $\{A', B\}$ of $\mathbb{D}$ which consists of smooth radial functions ρ_1 and ρ_2 such that

$$\max_i ||\nabla \rho_i||_{L^\infty(\mathbb{C})} \le \widetilde{C} w(B \cap A) < \widetilde{C}\epsilon$$

for some absolute constant $\widetilde{C} > 0$. Then using arguments similar to the above based on versions of (4.3), (4.4) and (4.5) for the cocycle c and the partition of unity $\{\rho_1, \rho_2\}$, we can find holomorphic functions $\overline{c}_1$ on B and $\overline{c}_2$ on A continuous on the corresponding boundaries such that

$$\overline{c}_1(z) - \overline{c}_2(z) = c(z), \quad z \in B \cap A, \quad \text{and}$$

$$\max\{||\overline{c}_1||_{H^\infty(B)}, ||\overline{c}_2||_{H^\infty(A)}\} \le \overline{C}||c||_{H^\infty(A\cap B)}$$

where $\overline{C} > 0$ is an absolute constant. Finally, define

$$F_\epsilon(z) := \begin{cases} f(z) - \overline{c}_1(z), & \text{if} \quad z \in B, \\ f_\epsilon(z) - \overline{c}_2(z), & \text{if} \quad z \in \mathbb{D} \setminus B. \end{cases}$$

Clearly we have

$$||f - F_\epsilon||_{L^\infty(\mathbb{D})} < c\epsilon.$$

for some absolute constant $c > 0$, and $F_\epsilon \in AP(F) \cap H^\infty$, where $F := \{z_1, \ldots, z_n\} \cap S$. Since ϵ is arbitrary, this completes the proof of the lemma. $\square$

As the next step of the proof let us establish Theorem 1.7 for $AP(F) \cap H^\infty$ with a finite set $F \subset \partial S$, say, $F = \{z_1, \ldots, z_n\}$.

Lemma 4.8

$$A_F = AP(F) \cap H^\infty.$$

Proof. Let us show that $AP(F) \cap H^\infty \subset A_F$. First suppose that F contains at least two points.

Let $\psi_1 : \partial\mathbb{D} \to \partial\mathbb{D}$ be the restriction to the boundary of a Möbius transformation of $\mathbb{D}$ that maps $-z_1$ to a point of F distinct from z_1 and preserves z_1. Then by the definition of Möbius transformations ψ_1 is a C^1 diffeomorphism of $\partial\mathbb{D}$. In particular, by Corollary 1.5 for a given $f \in AP(F) \cap H^\infty$ the function $f \circ \psi_1 \in AP(F_1) \cap H^\infty$ where $F_1 := \psi_1^{-1}(F)$. Since $z_1 \in F_1$, as in the proof of Theorem 4.5 we can find an almost periodic holomorphic function g_1 on $\overline{\mathbb{D}} \setminus \{\pm z_1\}$ such that $f \circ \psi_1 - g_1$ is continuous and equals 0 at z_1. We set

$$\widetilde{g}_1(z) := \frac{g(z)(z + z_1)}{2z_1}, \quad z \in \overline{\mathbb{D}} \setminus \{z_1\}.$$

Then $\widetilde{g}_1$ has a discontinuity at z_1 only. Let us show that $\widetilde{g}_1 \in A_{\{-z_1, z_1\}}$. Indeed, by the definition of g_1 the function $g_1 \circ \phi_{z_1}^{-1} \circ \mathrm{Log}^{-1}$ belongs to $AP_{\mathcal{O}}(\Sigma)$. Therefore, it can be uniformly approximated on Σ by polynomials in variables $e^{i\lambda z}$, $\lambda \in \mathbb{R}$, see, e.g., [JT]. In turn, g_1 can be uniformly approximated on $\overline{\mathbb{D}} \setminus \{\pm z_1\}$ by complex polynomials in variables $e^{i\lambda \mathrm{Log} \circ \phi_1}$. Now for $z \in \partial\mathbb{D}$ we have

$$\mathrm{Im}\{(\mathrm{Log} \circ \phi_{z_1})(z)\} := \begin{cases} 0, & \text{if} \quad 0 \le \mathrm{Arg}(z/z_1) < \pi, \\ \pi, & \text{if} \quad 0 \le \mathrm{Arg}(z_1/z) < \pi. \end{cases}$$

This implies that every function $e^{i\lambda \mathrm{Log}\circ\phi_1}$, $\lambda \in \mathbb{R}$, belongs to $A_{\{-z_1,z_1\}}$ and so $g_1 \in A_{\{-z_1,z_1\}}$, as well. Since $(z+z_1)/2z_1 \in A_0$, the function $\widetilde{g}_1 \in A_{\{-z_1,z_1\}}$ by definition. Thus the function $h_1 := \widetilde{g}_1 \circ \psi_1^{-1}$ belongs to A_F, is continuous outside z_1 and $f_1 := f - h_1 \in AP(F^1) \cap H^\infty$, where $F^1 := F \setminus \{z_1\}$. Further, using similar arguments we can find $h_2 \in A_F$, continuous outside z_2 such that $f_2 := f_1 - h_2 \in AP(F^2) \cap H^\infty$ where $F^2 := F^1 \setminus \{z_2\}$ etc. After n steps we obtain functions $h_1, \dots, h_n \in A_F$ such that h_k is continuous outside z_k, $1 \le k \le n$, and the function

$$h_{n+1} := f - \sum_{k=1}^{n} h_k \tag{4.6}$$

has no discontinuities on $\partial\mathbb{D}$, that is, $h_{n+1} \in A_0$. Therefore $f \in A_F$.

Next, if F consists of a single point, say z_0, then for a given $f \in AP(F) \cap H^\infty$ using the above argument we can find a function $h \in A_{\{z_0,z_1\}}$ with a fixed $z_1 \in \partial\mathbb{D}$ such that $f - h$ is continuous on F. Let $g \in A_0$ be a function equal to 1 at F and 0 at z_1. Then $f - gh \in A_0$. This completes the first part of the proof.

Show now that $A_F \subset AP(F) \cap H^\infty$. Again, assume first that F contains at least two points. Let $e^{\lambda f} \in A_F$, $\lambda \in \mathbb{R}$, where $\mathrm{Re}\, f$ is the characteristic function of an arc, say $[x,y]$, with $x, y \in F$. Let $\psi : \partial\mathbb{D} \to \partial\mathbb{D}$ be the restriction to $\partial\mathbb{D}$ of a Möbius transformation sending 1 to x and -1 to y. Then by the definition

$$(f \circ \psi \circ \phi_1^{-1})(z) = -\frac{i}{\pi}\mathrm{Log}\, z + C, \quad z \in \mathbb{H}_+,$$

for some constant C. Thus we have

$$e^{(\lambda f \circ \psi \circ (\mathrm{Log}\circ\psi_1)^{-1})(z)} = e^{\lambda C} e^{-i\lambda z/\pi}, \quad z \in \Sigma.$$

This means that $e^{\lambda f \circ \psi} \in AP(\{-1,1\}) \cap H^\infty$. Then by Corollary 1.5 we get $e^{\lambda f} \in AP(F) \cap H^\infty$. Since A_F is generated by A_0 and such functions $e^{\lambda f}$, we obtain the required implication.

If F is a single point, then we must show that $ge^{\lambda f} \in AP(F) \cap H^\infty$, $\lambda \in \mathbb{R}$, where $\mathrm{Re} f$ is the characteristic function of an arc with an endpoint at F and $g \in A_0$ is such that ge^f has discontinuity at F only. The result follows easily from the previous part of the proof, because $e^{\lambda f}$ is almost periodic on $\partial\mathbb{D} \setminus \{F, y\}$ for some y and is continuous at y.

This completes the proof of the lemma. $\square$

As it was mentioned above the required statement of the theorem follows from Lemmas 4.7 and 4.8 $\square$

4.3. Proof of Example 1.9. Using the bilinear transformation $\phi_1 : \mathbb{D} \to \mathbb{H}_+$, see (4.1), that maps 1 to $0 \in \mathbb{R}$ and -1 to ∞ we can transfer the problem to a similar one for functions on $\mathbb{H}_+$. Namely, let $\{x_k\}_{k\in\mathbb{N}} \subset \mathbb{R}_+$ be the sequence converging to 0, which is the image of the sequence $\{e^{it_k}\}_{k\in\mathbb{N}} \subset \partial\mathbb{D}$ of the example under ϕ_1. Let $H : \mathbb{R} \to \{0,1\}$ be the Heaviside function (i.e., the characteristic function of $[0,\infty)$). Then the pullback by ϕ_1^{-1} of the function u of the example to the boundary

$\mathbb{R}$ of $\mathbb{H}_+$ is the function

$$\widetilde{u}(x) := \sum_{k=1}^{\infty} \alpha_k H(x - x_k), \quad x \in \mathbb{R}.$$

We extend $\widetilde{u}$ to a harmonic function on $\mathbb{H}_+$ by the Poisson integral. Let $\widetilde{v}$ be the harmonic conjugate to the extended function determined on $\mathbb{R}$ by the formula

$$\widetilde{v}(x) = \sum_{k=1}^{\infty} \alpha_k \frac{\ln |x - x_k|}{\pi}, \quad x \in \mathbb{R}. \tag{4.7}$$

We set $\widetilde{h} := \widetilde{u} + i\widetilde{v}$. Then $\widetilde{h}$ is the pullback by ϕ_1^{-1} of a holomorphic function h on $\mathbb{D}$ such that $\operatorname{Re} h|_{\partial\mathbb{D}} = u$. Assume, to the contrary, that $e^h \in A_S$. Since $e^u \in R_S$, this assumption implies that $e^{iv} \in AP(S)$, where $v = \phi_1^*(\widetilde{v}|_{\mathbb{R}})$. Then according to the definition of the topology on $\mathcal{M}(AP(S))$, see section 3.3, the functions $\cos(\widetilde{v}(e^t))$ and $\sin(\widetilde{v}(e^t))$, $t \in \mathbb{R}$, admit continuous extensions to $b\mathbb{R}$ determined as follows.

If $\{s_\alpha\} \subset \mathbb{R}$ is a net converging in $b\mathbb{R}$ to a point $\eta \in b\mathbb{R}$, then the values at η of the extended functions are

$$\lim_{\alpha} \cos(\widetilde{v}(e^{s_\alpha})) \quad \text{and} \quad \lim_{\alpha} \sin(\widetilde{v}(e^{s_\alpha})),$$

respectively. In particular, this definition requires the existence of these limits.

Now, according to (4.7) there are sequences of points $\{x_k'\}_{k\in\mathbb{N}}$ and $\{x_k''\}_{k\in\mathbb{N}}$ in $\mathbb{R}_+$ such that x_k' and x_k'' are sufficiently close to x_k and

$$\lim_{k\to\infty} \left| \frac{x_k'}{x_k''} \right| = 1, \quad \cos(\widetilde{v}(x_k')) = 0, \quad \cos(\widetilde{v}(x_k'')) = 1, \quad k \in \mathbb{N}.$$

We set $t_k' := \ln x_k'$ and $t_k'' := \ln x_k''$, $k \in \mathbb{N}$. Assume without loss of generality that $\{t_k'\}$ forms a net converging in the topology of $b\mathbb{R}$ to a point $\xi \in b\mathbb{R}$ (for otherwise we replace $\{t_k\}$ by a proper subset satisfying this property). Since by the definition $\lim_{k\to\infty} |t_k' - t_k''| = 0$ and almost periodic functions on $\mathbb{R}$ are uniformly continuous, the family $\{t_k''\}$ forms a net with the same indeces as for the net formed by $\{t_k'\}$ whose limit in $b\mathbb{R}$ is ξ, as well. Hence, we must have

$$\lim_{k} \cos(\widetilde{v}(e^{t_k'})) = \lim_{k} \cos(\widetilde{v}(e^{t_k''})),$$

a contradiction. Therefore $e^h \notin A_S$.

Let now $f \in A_0$ be such that $f(1) = 0$. Then the function $(fe^h)|_{\partial\mathbb{D}}$ is continuous at 1. Thus $(fe^h)|_{\partial\mathbb{D}}$ can be uniformly approximated by constant functions on open arcs containing 1. The same is true for each point e^{it_k}. This implies immediately that $fe^h \in A_S$. $\square$

5. Proofs of Theorems 1.11 and 1.13.

5.1. Proof of Theorem 1.11. Let us recall that A_S is the uniform closure of the algebra generated by all possible A_F with finite $F \subset S$. Therefore the maximal ideal space $\mathcal{M}(A_S)$ of A_S is the inverse limit of the maximal ideal spaces $\mathcal{M}(A_F)$ of A_F. In particular, if we will prove that $\mathbb{D}$ is dense in each $\mathcal{M}(A_F)$, then by the definition of the inverse limit this will imply that $\mathbb{D}$ is dense in $\mathcal{M}(A_S)$, as required. Thus it suffices to prove the theorem for A_F with $F = \{z_1, \dots, z_n\} \subset \partial\mathbb{D}$.

Theorem 5.1 *$\mathbb{D}$ is dense in $\mathcal{M}(A_F)$.*

Proof. Let $\mathcal{I}_k \subset A_F$ be the closed ideal consisting of functions that are continuous and equal to 0 at z_k. By A_k we denote the quotient Banach algebra $A_F/\mathcal{I}_k$ equipped with the quotient norm. Let us recall that $\mathcal{M}_{z_k}$ is the maximal ideal space of the algebra $\mathcal{A}_{z_k}$ which is the uniform closure of the algebra of continuous functions on $\mathbb{D}$ almost periodic in circular neighbourhoods of z_k. Also, according to Lemma 4.4 (b) there is a natural continuous projection $p_{z_k} : \mathcal{M}_{z_k} \to \overline{\mathbb{D}}$ and $p_{z_k}^{-1}(z_k)$ is homeomorphic to $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$. Moreover, by Lemma 4.4 (c) each $f \in A_F$ is extended to a continuous function on $\mathcal{M}_{z_k}$ holomorphic on $p_{z_k}^{-1}(z_k)$. Hence, there is a continuous map $H_k : \mathcal{M}_{z_k} \to \mathcal{M}(A_F)$ whose image coincides with the closure of $\mathbb{D}$. Moreover, according to the decomposition obtained in the proof of Lemma 4.8, see (4.6), H_k maps $p_{z_k}^{-1}(z_k)$ homeomorphically onto its image.

Lemma 5.2 *Let $\phi_k : A_F \to AP_{\mathcal{O}}(\Sigma)$ be the composition of the extension homomorphism of functions from A_F to $\mathcal{M}_{z_k}$ and the restriction homomorphism of functions on $\mathcal{M}_{z_k}$ to $p_{z_k}^{-1}(z_k)$. Then $Ker\,\phi_k = \mathcal{I}_k$ and A_k is isomorphic to $AP_{\mathcal{O}}(\Sigma)$.*

Proof. Clearly, $\mathcal{I}_k \subset Ker\ \phi_k$. Let us check the converse implication. Let $f \in Ker\ \phi_k$. As it follows from the proof of Lemma 4.8, see (4.6), there are linear continuous operators $T_k : AP_{\mathcal{O}}(\Sigma) \to A_{\{z_k\}} \subset A_F$ such that $\phi_k \circ T_k = id$. Moreover, $T_0 := I - \sum_{k=1}^{n} T_k \circ \phi_k$, where I is the identity map, maps A_F onto A_0. In particular, we have $T_k(\phi_k(f)) = 0$. Thus $f = -T_0(f) + \sum_{s \neq k} T_s(\phi_s(f))$. Since $\phi_k(f) = 0$, this implies that f is continuous and equal to 0 at z_k. Now from the formula $\phi_k \circ T_k = id$ we obtain that A_k is isomorphic to $AP_{\mathcal{O}}(\Sigma)$. $\square$

Let $i : A_0 \hookrightarrow A_F$ be the natural inclusion. Its dual determines a continuous surjective map $a_F : \mathcal{M}(A_F) \to \overline{\mathbb{D}}$. Next, taking the dual map to ϕ_k we obtain that each $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ is embedded into $\mathcal{M}(A_F)$, its image coincides with $H_k(p_{z_k}^{-1}(z_k))$ and a_F maps $H_k(p_{z_k}^{-1}(z_k))$ to z_k.

Let $\xi \in \mathcal{M}(A_F)$ and $\mathfrak{m} := \text{Ker}\ \xi \subset A_F$ be the corresponding maximal ideal. Assume first that there is k such that $\mathcal{I}_k \subset \mathfrak{m}$. Then $\mathfrak{m}_k = \phi_k(\mathfrak{m})$ is a maximal ideal of A_k. By $\xi_k \in \mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ we denote the character corresponding to $\mathfrak{m}_k$. Then $\xi = \phi^*(\xi_k) \in H_k(p_{z_k}^{-1}(z_k))$. Now, by the definition of H_k the point ξ belongs to the closure of $\mathbb{D}$ in $\mathcal{M}(A_F)$. We continue with the following lemma.

Lemma 5.3 *Assume that a maximal ideal $\mathfrak{m}$ of A_F does not contain any of $\mathcal{I}_k$. Then $\mathfrak{m}$ does not contain $\cap_{1 \le k \le n} \mathcal{I}_k$, as well.*

Proof. Suppose, to the contrary, that $\cap_{1 \le k \le n} \mathcal{I}_k \subset \mathfrak{m}$. Let $x_k \in \mathcal{I}_k$, $1 \le k \le n$, be such that $x_k \notin \mathfrak{m}$. Since $\mathcal{I}_k$ are ideals, $x_1 \cdots x_n \in \cap_{1 \le k \le n} \mathcal{I}_k$. Thus $x_1 \cdots x_n \in \mathfrak{m}$. Since $\mathfrak{m}$ is a prime ideal, there is some k so that $x_k \in \mathfrak{m}$, a contradiction. $\square$

From this lemma it follows that in order to prove the theorem it remains to consider the case $\mathfrak{m} \not\in \cap_{1 \le k \le n} \mathcal{I}_k$. Observe that $\cap_{1 \le k \le n} \mathcal{I}_k$ consists of all functions from A_0 that vanish on F. Thus there is $f \in \cap_{1\leq k\leq n}\mathcal{I}_k$ such that $f(\xi) \neq 0$. This implies that $a_F(\xi) \notin F$. For every $g \in A_F$, let us consider the function gf. By the definition $gf \in A_0$. Thus we have

$$g(a_F(\xi))f(a_F(\xi)) = (gf)(a_F(\xi)) = (gf)(\xi) = g(\xi)f(\xi) = g(\xi)f(a_F(\xi)).$$

Equivalently,

$$g(\xi) = g(a_F(\xi)) \quad \text{for all} \quad g \in A_F.$$

This implies that $a_F^{-1}(a_F(\xi)) = \{\xi\}$. Therefore $a_F : \mathcal{M}(A_F) \setminus a_F^{-1}(F) \to \overline{\mathbb{D}} \setminus F$ is a homeomorphism. In particular, ξ belongs to the closure of $\mathbb{D}$.

The proof of Theorem 1.11 is complete. □

5.2. Proof of Theorem 1.13. Statements (1) and (3) of the theorem follow easily from similar statements for a_F with a finite subset $F \subset S$ proved in section 5.1 and the properties of the inverse limit. Let us prove (3). We first prove the statement for a finite subset $F \subset S$.

Since $A_0 \subset A_F$ and each function from A_0 attains the maximum of modulus on $\partial\mathbb{D}$, $K_F \subset a_F^{-1}(\partial\mathbb{D})$. As it follows from Theorem 1.7 $A_F \hookrightarrow AP(F)$. Also, the extensions of functions from A_F to $\mathcal{M}(AP(F))$ separate points there. Therefore $\mathcal{M}(AP(F))$ is embedded into $\mathcal{M}(A_F)$. Identifying $\mathcal{M}(AP(F))$ with its image under this embedding we have $\mathcal{M}(AP(F)) \subset a_F^{-1}(\partial\mathbb{D})$. Since $a_F^{-1}(\partial\mathbb{D}) \setminus a_F^{-1}(F) \to \partial\mathbb{D} \setminus F$ is a homeomorphism and each $z \in \partial\mathbb{D}$ is a peak point for A_0, the set K_F contains the closure of $\mathbb{D} \setminus F$ which is, by definition, coincides with $\mathcal{M}(AP(F))$. Assume that there is $\xi \in K_F \setminus \mathcal{M}(AP(F))$. Then $a_F(\xi) := z^* \in F$. Further, identifying $a_S^{-1}(z^*)$ with $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$ we get from our assumption that $\xi \in i_\eta(\Sigma_0)$ for some $\eta \in b\mathbb{Z}$, see section 2; here Σ_0 is the interior of Σ. Then, because $i_\eta(\Sigma_0)$ is dense in $\mathcal{M}(AP_{\mathcal{O}}(\Sigma))$, by the maximum modulus principle each function $f \in A_F$ for which $\max_{\mathbb{D}} |f| = |f(\xi)|$ is constant on $a_S^{-1}(z^*)$. Thus such f admits the maximum of modulus on $\mathcal{M}(AP(F))$ also. This contradicts to the minimality of K_F. Therefore $K_F = \mathcal{M}(AP(F))$.

Further, $\mathcal{M}(AP(S))$ is the inverse limit of compact sets $\mathcal{M}(AP(F))$ for all finite $F \subset S$. As before we naturally identify $\mathcal{M}(AP(S))$ with a subset of $\mathcal{M}(A_S)$. Then since A_S is the uniform closure of the algebra generated by all possible A_F with finite $F \subset S$, by the definition of the inverse limit $K_S \subseteq \mathcal{M}(AP(S))$. But in fact K_S coincides with $\mathcal{M}(AP(S))$ because otherwise its projection to some of $\mathcal{M}(A_F)$ is a boundary of A_F and a proper subset of $\mathcal{M}(AP(F))$, a contradiction. □

References

[B] H. Bohr, Almost periodic functions. Chelsea, New York, 1947.

[Br] A. Brudnyi, Topology of the maximal ideal space of H^∞, J. Funct. Anal., **189** (1) (2002), 21-52.

[Bö] A. Bötcher, On the corona theorem for almost periodic functions. Integral Equations Operator Theory **33** (3) (1999), 253-272.

[D] J. Dieudonne, Foundations of modern analysis, Academic Press, New York, 1969.

[G] J. Garnett, Bounded analytic functions. Academic Press, New York, 1981.

[H] F. Hirzebruch, Topological methods in Algebraic Geometry. Springer-Verlag, New York, 1966.

[JT] B. Jessen and H. Tornehave, Mean motions and zeros of almost periodic functions. Acta Math. **77** (1945), 137-279.

[L] E. Lindelöf, Sur une principe générale de l'analyse et ces applications à la théorie de la representation conforme. Acta Soc. Sci. Fenn. **46** (4) (1915).

[LZ] B.M. Levitan and V.V. Zhikov, Almost periodic functions and differential equations. Cambridge University Press, 1982.

[P] L. S. Pontryagin, Selected works. Vol. 2. Gordon & Beach Science Publishers, New York, 1986.

[R] H. Royden, Function algebras, Bull. Amer. Math. Soc. **69** (1963), 281-298.